

Borel summability and Lindstedt series

O. Costin¹, G. Gallavotti², G. Gentile³ and A. Giuliani⁴

¹ Department of Mathematics, Rutgers University

² Dipartimento di Fisica, Università di Roma Uno, I-00185

³ Dipartimento di Matematica, Università di Roma Tre, Roma, I-00146

⁴ Dipartimento di Matematica, Università di Roma Due, Roma, I-00133

ABSTRACT. *Resonant motions of integrable systems subject to perturbations may continue to exist and to cover surfaces with parametric equations admitting a formal power expansion in the strength of the perturbation. Such series may be, sometimes, summed via suitable sum rules defining C^∞ functions of the perturbation strength: here we find sufficient conditions for the Borel summability of their sums in the case of two-dimensional rotation vectors with Diophantine exponent $\tau = 1$ (e.g. with ratio of the two independent frequencies equal to the golden mean).*

sec.1

1. Introduction

In the paradigmatic setting of KAM theory, one considers unperturbed motions $\varphi \rightarrow \varphi + \omega_0(\mathbf{I})t$ on the torus \mathbb{T}^d , $d \geq 2$, driven by a Hamiltonian $H = H_0(\mathbf{I})$, where $\mathbf{I} \in \mathbb{R}^d$ are the actions conjugated to φ and $\omega_0(\mathbf{I}) = \partial_{\mathbf{I}} H_0(\mathbf{I})$. Standard analytic KAM theorem considers the perturbed Hamiltonian $H_\varepsilon = H_0(\mathbf{I}) + \varepsilon f(\varphi, \mathbf{I})$, with f analytic, and a frequency vector ω_0 which is Diophantine with constants C_0 and τ (i.e. $|\omega_0 \cdot \nu| \geq C_0 |\nu|^{-\tau} \forall \nu \in \mathbb{Z}^d, \nu \neq \mathbf{0}$) and which is among the frequencies of the unperturbed system: $\omega_0 = \omega_0(\mathbf{I}_0)$ for some \mathbf{I}_0 . Suppose $H_0(\mathbf{I}) = \mathbf{I}^2/2$ and $f(\varphi, \mathbf{I}) = f(\varphi)$ for simplicity, that is assume that H_0 is quadratic and the perturbation depends only on the angle variables. Then for ε small enough the unperturbed motion $\varphi \rightarrow \varphi + \omega_0 t$ can be analytically continued into a motion of the perturbed system, in the sense that there is an ε -analytic function $\mathbf{h}_\varepsilon : \mathbb{T}^d \rightarrow \mathbb{T}^d$, reducing to the identity as $\varepsilon \rightarrow 0$, such that $\psi + \mathbf{h}_\varepsilon(\psi)$ solves the Hamilton equations for H_ε , i.e. $\ddot{\varphi} + \varepsilon \partial_\varphi f(\varphi) = \mathbf{0}$, if ψ is replaced by $\psi + \omega t$, for any choice of the initial data ψ . The function \mathbf{h}_ε (called the conjugation) can be constructed as a power series in ε (Lindstedt series) and for ε small convergence can be proved, exploiting cancellations and summation methods typical of quantum field theory, [GBG]. We can call this the *maximal* KAM theorem, as it deals with invariant tori of maximal dimension.

The same methods allow us to study existence of perturbed resonant quasi-periodic motions in quasi-integrable Hamiltonian systems. By resonant here we mean that ω_0 satisfies $1 \leq s \leq d - 1$ rational relations, i.e. it can be reduced to a vector $(\omega, \mathbf{0})$, $\omega \in \mathbb{R}^{d-s}$, with rationally-independent components, via a canonical transformation acting as a linear integer coefficients map of the angles. In this representation we shall write $\varphi = (\alpha, \beta)$ denoting by α the “fast variables” rotating with angular velocity ω and by β the fixed unperturbed angles (“slow variables”).

The study of resonant quasi-periodic motions is mathematically a natural extension of the maximal KAM case and physically it arises in several stability problems. We mention here celestial mechanics, where the phenomenon of resonance locking between rotation and orbital periods of satellites is a simple example (as in this case the resonant torus is one-dimensional, i.e. it describes a periodic motion). In general resonant motions arise in presence of small friction: the most unstable motions are the maximally quasi-periodic ones (on KAM tori). In presence of friction the maximally quasi-periodic motions “collapse” into resonant motions with one frequency rationally related to the others, then on a longer time scale one more frequency gets locked to the others and the motion takes place on an invariant torus with dimension lower than the maximal by 2, and so on. Periodic motions are the least dissipative, and eventually the motion becomes maximally resonant, i.e. periodic. This is a scenario among others possible, and its study in particular cases seems to require a good understanding of the properties of the resonant motions of any dimension.

The first mathematical result is that also in the resonant case, if ω is a $(d - s)$ -dimensional Diophantine vector, a conjugation \mathbf{h}_ε can be constructed by summations of the Lindstedt series; however the conjugation \mathbf{h}_ε that one is able to construct is not analytic, but, at best, only C^∞ in ε : in general it is defined only on a large measure Cantor set \mathcal{E} of ε 's, $\mathcal{E} \subset [-\varepsilon_0, \varepsilon_0]$ for some positive ε_0 , so that C^∞ has to be meant in the sense of Whitney. It is commonly believed that a conjugation which is analytic in a domain including the origin does not exist in the resonant case. Moreover, contrary to what happens in the maximal case, not all resonant unperturbed motions with a given rotation vector ω appear to survive under perturbation, but only a discrete number of them. This is not due to technical limitations of the method, and it has the physical meaning that only points β_0 which are equilibria for the “effective potential” $(2\pi)^{s-d} \int d\alpha f(\alpha, \beta)$ can remain in average at rest in presence of the perturbation.

The following natural (informal) question then arises: “where do the unperturbed motions corresponding to initial data (α, β) , $\beta \neq \beta_0$, disappear when we switch on the perturbation?” An intriguing scenario is that the tori that seem to disappear in the construction of \mathbf{h}_ε actually “condense” into a continuum of highly degenerate tori near the ones corresponding to the equilibria β_0 . It is therefore interesting to study “uniqueness”, regularity and possible degeneracies of the perturbed tori constructed by the Lindstedt series methods (or by alternative methods, such as classical Newton’s iteration scheme).

In the present paper we investigate *hyperbolic* (see below) perturbed tori with two-dimensional rotation vectors for a class of analytic quasi-integrable Hamiltonians. Informally, our main result is that the hyperbolic tori, which survive to the switching of the perturbation and are described by a function \mathbf{h}_ε that we construct explicitly, are independent on the procedure used to construct them iteratively (which is not obvious, due to lack of analyticity). Moreover, if $\eta = \sqrt{\varepsilon}$, *the conjugation is Borel summable* in η for $\varepsilon > 0$. *The conjugation constructed here is the unique possible for our problem within the class of Borel summable functions.* Of course this does not exclude existence of other less regular quasi-periodic motions, not even existence of other quasi-periodic solutions which admit the same formal power expansion as \mathbf{h}_ε .

In order to make our results more precise, we first introduce the model and summarize the results about existence and properties of lower-dimensional perturbed tori as we need in the following. Then we briefly recall the definition and some key properties of Borel summable functions, and finally we state more technically our main result.

p.1.1 **1.1. The model.**

Consider a Hamiltonian system

$$1.0 \quad H = \frac{1}{2}\mathbf{A}^2 + \frac{1}{2}\mathbf{B}^2 + \eta^2 f(\boldsymbol{\alpha}, \boldsymbol{\beta}), \quad (1.1)$$

with $\mathbf{I} = (\mathbf{A}, \mathbf{B}) \in \mathbb{R}^r \times \mathbb{R}^s$ the action coordinates and $\boldsymbol{\varphi} = (\boldsymbol{\alpha}, \boldsymbol{\beta}) \in \mathbb{T}^r \times \mathbb{T}^s$ the conjugated angle coordinates. Let $\mathbf{A}_0 = \boldsymbol{\omega}, \mathbf{B} = \mathbf{0}$ be an unperturbed resonance with $|\boldsymbol{\omega} \cdot \boldsymbol{\nu}| \geq C_0 |\boldsymbol{\nu}|^{-\tau}$ for $\boldsymbol{\nu} \in \mathbb{Z}^r$, $\boldsymbol{\nu} \neq \mathbf{0}$ and for some $C_0, \tau > 0$, and let $((\mathbf{A}_0, \mathbf{0}), (\boldsymbol{\psi} + \boldsymbol{\omega} t, \boldsymbol{\beta}_0))$ be the corresponding unperturbed resonant motions with initial angles $\boldsymbol{\alpha} = \boldsymbol{\psi}, \boldsymbol{\beta} = \boldsymbol{\beta}_0$.

The following result holds.

Proposition 1. ([GG1],[GG2]) *Let $\boldsymbol{\beta}_0$ be a non-degenerate maximum of the function $f_0(\boldsymbol{\beta}) \equiv (2\pi)^{-r} \int d\boldsymbol{\alpha} f(\boldsymbol{\alpha}, \boldsymbol{\beta})$, i.e. $\partial_{\boldsymbol{\beta}} f_0(\boldsymbol{\beta}_0) = 0$ and $\partial_{\boldsymbol{\beta}}^2 f_0(\boldsymbol{\beta}_0) < 0$, and let $\boldsymbol{\omega} \in \mathbb{R}^r$ be a Diophantine vector of constants C_0, τ , i.e. $|\boldsymbol{\omega} \cdot \boldsymbol{\nu}| \geq C_0 |\boldsymbol{\nu}|^{-\tau}$ for all $\boldsymbol{\nu} \in \mathbb{Z}^r$, $\boldsymbol{\nu} \neq \mathbf{0}$. Then for $\varepsilon > 0$, setting $\eta = \sqrt{\varepsilon}$, there exists a function $\boldsymbol{\psi} \rightarrow \mathbf{h}(\boldsymbol{\psi}, \eta) = (\mathbf{a}(\boldsymbol{\psi}), \mathbf{b}(\boldsymbol{\psi}))$, vanishing as $\eta \rightarrow 0$ and with the following properties.*

(i) *The functions $t \rightarrow \boldsymbol{\varphi}(t) = (\boldsymbol{\alpha}(t), \boldsymbol{\beta}(t)) = (\boldsymbol{\psi} + \boldsymbol{\omega} t, \boldsymbol{\beta}_0) + \mathbf{h}(\boldsymbol{\psi} + \boldsymbol{\omega} t, \eta)$ satisfy the equation of motion $\ddot{\boldsymbol{\varphi}} = -\varepsilon \partial_{\boldsymbol{\varphi}} f(\boldsymbol{\varphi})$ for any choice of $\boldsymbol{\psi} \in \mathbb{T}^r$.*

(ii) *The function \mathbf{h} is defined for $(\boldsymbol{\psi}, \eta) \in \mathbb{T}^r \times [0, \eta_0]$ and can be analytically continued to an holomorphic function in the domain $\mathcal{D} = \{\eta \in \mathbb{C} : \text{Re } \eta^{-1} > \eta_0^{-1}\}$.*

(iii) *(The analytic continuation of) \mathbf{h} is C^∞ in η at the origin along any path contained in \mathcal{D} and its Taylor coefficients at the origin satisfy, for suitable positive constants C and D , the bounds $|\frac{1}{k!} \partial_\eta^k \mathbf{h}(\boldsymbol{\psi}, 0)| < DC^k k!^\tau$, where τ is the Diophantine exponent of $\boldsymbol{\omega}$.*

Remarks.

(1) The domain \mathcal{D} is a circle in \mathbb{C}^2 , centered in $\eta_0/2$ and of radius $\eta_0/2$ (hence tangent to the imaginary axis at the origin).

(2) A function \mathbf{h} was constructed in [GG1] by a perturbative expansion in power series in $\varepsilon = \eta^2$ (Lindstedt series) and by exploiting multiscale decomposition, cancellations and summations in order to control convergence of the series. Eventually \mathbf{h} is expressed as a new series which is not a power series in $\varepsilon = \eta^2$ and which is holomorphic in \mathcal{D} . The procedure leading to the convergent resummed series from the initial formal power series in $\varepsilon = \eta^2$ relies on a number of arbitrary choices (which will be made explicit in next section) and a priori is not clear that the result is actually independent on such choices. Another (in principle) function \mathbf{h} was constructed in [GG2], with a method which applies in more general cases (see item (4) below): we shall see that the two functions in fact coincide.

(3) The function $\mathbf{h}(\boldsymbol{\psi}, \eta)$ can be regarded as a function of $\varepsilon = \eta^2$, as in [GG1] and [GG2]. As such the bound in item (iii) would be modified into $|\frac{1}{k!} \partial_\varepsilon^k \mathbf{h}(\boldsymbol{\psi}, 0)| < DC^k (2k)!^\tau$, and the analyticity domain in item (ii) would be as described in [GG1], Fig. 1. We also note here that the exponent $2\tau + 1$ in [GG1], Eq. (5.29), was not correct (without consequences as the value of the exponent was just quoted and not exploited), as the right one is 3τ : of course for $\tau = 1$ (that is the case we consider in this paper) the two values coincide.

(4) In [GG2] a similar statement was proved for $\boldsymbol{\beta}_0$ a non-degenerate equilibrium point of the function $f_0(\boldsymbol{\beta})$, i.e. $\boldsymbol{\beta}_0$ not necessarily a maximum. If $\boldsymbol{\beta}_0$ is a maximum the corresponding torus is called *hyperbolic*, if $\boldsymbol{\beta}_0$ is a minimum it is called *elliptic*. In the elliptic case for ε real the domain of definition of \mathbf{h} on the real line is $\mathbb{T}^r \times \mathcal{E}$, where $\mathcal{E} \subset [0, \varepsilon_0]$ is a set with open dense complement in $[0, \varepsilon_0]$ but with 0 as a density point in the sense of Lebesgue. The reason for stating Proposition

1 as above is that *in the present paper we shall restrict our analysis to the hyperbolic case.*

(5) If $\tau = 1$, i.e. if $r = 2$ and ω is quadratically irrational, then the bound on the coefficients of the power expansion of h in η at the origin is $|\mathbf{h}^{(k)}(\psi)| < DC^k k!$. Hence in this case it is natural to ask for Borel summability of \mathbf{h} .

(6) Even if we consider only hyperbolic tori, in the following we shall use the method introduced in [GG2], because it is more general and it is that one should look at if one tried to extend the analysis to the case of elliptic tori. The main difference between the forthcoming analysis and that of [GG2] is the use of a sharp multiscale decomposition, instead of a smooth one, as it allows further simplifications in the case of hyperbolic tori. We shall come back to this later.

p.1.2 **1.2. Borel transforms.** Let $F(\eta)$ be a function of η which is analytic in a disk centered at $(\frac{1}{2\rho_0}, 0)$ and radius $\frac{1}{2\rho_0}$ (i.e. centered on the positive real axis and tangent, at the origin, to the imaginary axis), and which vanishes as $\eta \rightarrow 0$ as η^q for some $q > 1$. Then one can consider the inverse Laplace transform of the function $z \rightarrow F(z^{-1})$, defined for p real and positive and $\rho > \rho_0$ by

$$2.1 \quad \mathcal{L}^{-1}F(p) = \int_{\rho-i\infty}^{\rho+i\infty} e^{z p} F\left(\frac{1}{z}\right) \frac{dz}{2\pi i}. \quad (1.2)$$

If F admits a Taylor series at the origin in the form $F(\eta) \sim \sum_{k=2}^{\infty} F_k \eta^k$ then the Taylor series for $\mathcal{L}^{-1}F$ at the origin is

$$2.2 \quad \mathcal{L}^{-1}F(p) \sim \sum_{k=2}^{\infty} F_k \frac{p^{k-1}}{(k-1)!}. \quad (1.3)$$

If the series in (1.3) is convergent then the sum of the series coincides with $\mathcal{L}^{-1}F(p)$ for $p > 0$ real. Of course the series expansion (1.3) provides an expression suitable for studying analytic continuation of $\mathcal{L}^{-1}F(p)$ outside the real positive axis. Note that (1.3) makes sense even in the case the series starts from $k = 1$. Then given any formal power series $F(\eta) \sim \sum_{k=1}^{\infty} F_k \eta^k$ we shall define $F_B(p) = \sum_{k=1}^{\infty} F_k \frac{p^{k-1}}{(k-1)!}$ as the *Borel transform* of $F(\eta)$, whenever the sum defining it is convergent. It is remarkable that in some cases the map $F \rightarrow F_B$ is invertible. If this is the case one says that the Taylor series of F is *Borel summable* and we also say that the function F is *Borel summable*: this is made precise as follows.

Definition 1. Let a function $\eta \rightarrow F(\eta)$ be such that

(i) it is analytic in a disk centered at $(\frac{1}{2\rho_0}, 0)$ and radius $\frac{1}{2\rho_0}$, and admits an asymptotic Taylor series at the origin where it vanishes,

(ii) its Taylor series at the origin admits a Borel transform $F_B(p)$ which is analytic for p in a neighborhood of the positive real axis, and on the positive real axis grows at most exponentially as $p \rightarrow +\infty$,

(iii) it can be expressed, for $\eta > 0$ small enough, as

$$2.3 \quad F(\eta) = \int_0^{+\infty} e^{-p/\eta} F_B(p) dp. \quad (1.4)$$

Then we call F Borel summable (at the origin).

Remarks.

(1) If F is Borel summable, then one says that F is equal to the Borel sum of its own Taylor series.

(2) For instance one checks that a function F holomorphic at the origin and vanishing at the origin is Borel summable; its Borel transform is entire.

(3) The function $F(\eta) = \sum_{k=1}^{\infty} 2^{-k} \frac{\eta}{1+k\eta}$ is not analytic at the origin but it is Borel summable.

(4) If F, G are Borel summable then also FG is Borel summable and $(FG)_B(p) = \int_0^p F_B(p')G_B(p-p')dp' \equiv (F_B * G_B)(p)$ on the common analyticity domain of F_B and G_B , with the integral which can be computed along any path from 0 to p in the common analyticity domain. Of course by definition $|F_B * G_B| \leq |F_B| * |G_B|$, where in the r.h.s. the convolution is along any path from 0 to p in the common analyticity domain of F_B and G_B (note however that now the convolution $|F_B| * |G_B|$ depends on the path).

(5) The Borel transform of η^k is $p^{k-1}/(k-1)!$. The Borel transform of $\eta/(1-\alpha\eta)$ is $e^{\alpha p}$. If $F_B(p) = e^{\alpha p} p^{k_1}/k_1!$ and $G_B(p) = e^{\beta p} p^{k_2}/k_2!$, then $F_B * G_B(p) = e^{\alpha p} p^{k_1+k_2+1}/(k_1+k_2+1)!$

(6) More generally if F_i , $i = 1, 2$, have Borel transforms F_{iB} bounded in a sector around the real axis and centered at the origin by $|F_{iB}(p)| \leq C e^{\alpha_i |p| + \beta_i |\text{Im } p|} |p|^{k_i}/k_i!$, with α_i and β_i real, then $|F_{1B} * F_{2B}(p)| \leq C^2 e^{\alpha |p| + \beta |\text{Im } p|} |p|^{k_1+k_2+1}/(k_1+k_2+1)!$ where $\alpha = \max \alpha_i$ and $\beta = \max \beta_i$. We shall make use of this bound repeatedly below.

p.1.3

1.3. Main results. We are now ready to state more precisely our main results.

Proposition 2. *Let us consider Hamiltonian (1.1) with $r = 2$ and $f(\alpha, \beta)$ an analytic function of its arguments. Let $\beta_0 \in \mathbb{T}^s$ satisfy the assumptions of Proposition 1 and $\omega \in \mathbb{R}^2$ be a Diophantine vector with constants $C_0 > 0$ and $\tau = 1$, i.e. $|\omega \cdot \nu| \geq C_0 |\nu|^{-1}$ for all $\nu \in \mathbb{Z}^T$, $\nu \neq \mathbf{0}$. Then there exists a unique Borel summable function $\mathbf{h}(\psi, \eta)$ satisfying properties (i)–(iii) of Proposition 1.*

Note that, once existence of a Borel summable function $\mathbf{h}(\psi, \eta)$ satisfying properties (i)–(iii) of Proposition 1 is obtained, the uniqueness in the class of Borel summable functions is obvious, by the very definition of Borel summability. In fact all such functions have a Borel transform $\mathbf{h}_B(\psi, p)$ that is p -analytic in an open domain enclosing \mathbb{R}^+ (hence also in a neighborhood of the origin), where they coincide (because they all have the same expansion at the origin), then they all coincide everywhere.

The proof of Proposition 2 will proceed by showing Borel summability of (one of) the function(s) $\mathbf{h}(\psi, \eta)$ constructed in [GG2]. In particular a corollary of the proof is that $\mathbf{h}(\psi, \eta)$ constructed in [GG2] is independent of the arbitrary choices mentioned in Remark (2) after Proposition 1.

Our proof of Borel summability of \mathbf{h} does not use Nevanlinna's theorem, [Ne] [So]. In fact we failed in checking that \mathbf{h} satisfies the hypothesis of the theorem. Our strategy goes as follows. We introduce a sequence of approximants $\mathbf{h}^{(N)}$ to \mathbf{h} , that is naturally induced by the multiscale construction of [GG2]. We explicitly check that $\mathbf{h}^{(1)}$ is Borel summable and that its Borel transform is entire. Then we show inductively that the analyticity domain of $\mathbf{h}_B^{(N)}$ is a neighborhood \mathcal{B} of \mathbb{R}^+ (not shrinking to 0 as $N \rightarrow \infty$) and that $\mathbf{h}_B^{(N)}$ grows very fast at infinity in \mathcal{B} (in general faster than exponential). However the results of Proposition 1 imply that the growth of $\mathbf{h}_B^{(N)}$ on the positive real line is uniformly bounded by an exponential. Then Borel summability of \mathbf{h} follows by performing the limit $N \rightarrow \infty$ and using uniform bounds that we shall derive on the approximants and on their Borel transforms.

In the next section we will recall the structure and the properties of the resummed series obtained in [GG2], defining the function $\mathbf{h}(\psi, \eta)$ of Proposition 1. In Section 3 we define the sequence $\mathbf{h}^{(N)}$ of approximants and we show that the inverse Laplace transform of $\mathbf{h}^{(N)}$ is uniformly bounded by an exponential on the positive real line. In Section 4 we prove Borel summability of $\mathbf{h}(\psi, \eta)$ in

the easier case in which the perturbation $f(\boldsymbol{\alpha}, \boldsymbol{\beta})$ in (1.1) is a trigonometric polynomial in $\boldsymbol{\alpha}$. In Appendix A1 we discuss how to extend the method to cover the general analytic case. Finally, in Appendix A2 we show that the same result applies to the function \mathbf{h} constructed in [GG1]: this allows us to identify the functions constructed with the two methods of [GG1] and [GG2], since they are both Borel summable and admit the same formal expansion at the origin.

sec.3

2. Lindstedt series

Denote by $\mathbf{a}(\boldsymbol{\psi})$, $\mathbf{b}(\boldsymbol{\psi})$ the $\boldsymbol{\alpha}$, $\boldsymbol{\beta}$ components of \mathbf{h} , respectively. In [GG2] an algorithm is described to construct order by order in η^2 the solution to the homologic equation

$$\begin{cases} (\boldsymbol{\omega} \cdot \partial_{\boldsymbol{\psi}})^2 \mathbf{a}(\boldsymbol{\psi}) = \eta^2 \partial_{\boldsymbol{\alpha}} f(\boldsymbol{\psi} + \mathbf{a}(\boldsymbol{\psi}), \boldsymbol{\beta}_0 + \mathbf{b}(\boldsymbol{\psi})), \\ (\boldsymbol{\omega} \cdot \partial_{\boldsymbol{\psi}})^2 \mathbf{b}(\boldsymbol{\psi}) = \eta^2 \partial_{\boldsymbol{\beta}} f(\boldsymbol{\psi} + \mathbf{a}(\boldsymbol{\psi}), \boldsymbol{\beta}_0 + \mathbf{b}(\boldsymbol{\psi})). \end{cases} \quad (2.1)$$

The resulting series, called the ‘‘Lindstedt series’’, is widely believed to be divergent. A summation procedure has been found which collects its terms into families until a convergent series is obtained. The resummed series (no longer a power series) can be described in terms of suitably decorated tree graphs, i.e. \mathbf{h} can be expressed as a sum of values of tree graphs:

$$h_{\gamma, \boldsymbol{\nu}}(\eta) = \sum_{\theta \in \Theta_{\boldsymbol{\nu}, \gamma}} \text{Val}(\theta), \quad (2.2)$$

where $\mathbf{h}_{\boldsymbol{\nu}}$ is the $\boldsymbol{\nu}$ -th coefficient in the Fourier series for \mathbf{h} , and $\gamma = \{1, \dots, 2 + s\}$ labels the component of the vector $\mathbf{h}_{\boldsymbol{\nu}}$ (recall that $2 + s$ is the number of degrees of freedom of our Hamiltonian, 2 being the number of ‘‘fast variables’’ $\boldsymbol{\alpha}$ and s being the number of ‘‘slow variables’’ $\boldsymbol{\beta}$). $\Theta_{\boldsymbol{\nu}, \gamma}$ is the set of decorated trees contributing to $h_{\gamma, \boldsymbol{\nu}}(\eta)$ and, given $\theta \in \Theta_{\boldsymbol{\nu}, \gamma}$, $\text{Val}(\theta)$ is its value, both still to be defined.




FIGURE 1. A tree θ with 12 nodes; one has $p_{v_0} = 2, p_{v_1} = 2, p_{v_2} = 3, p_{v_3} = 2, p_{v_4} = 2$. The length of the lines should be the same but it is drawn of arbitrary size. The nodes $v_i, i = 5, \dots, 11$ will be called endnodes. The separated line illustrates the way to think of the label $\eta = (\gamma', \gamma)$.

We now describe the rules to construct the tree graphs and to compute their value. We shall need the explicit structure in the proof of Borel summability in next sections, and this is why we are reviewing it here. Given the rules below one can formally check that the sum (2.2) is a solution to the Hamilton equations, see [GG2]. A few differences (in fact simplifications) arise

here with respect to [GG2], and we provide some details with the aim of making the discussion self-consistent. Essentially, the changes consist of: (i) shifting the order of factors in products appearing in the definition of the values $\text{Val}(\theta)$ to an order that makes it easier to organize the recursive evaluation of several Borel transforms; see remarks following (2.9), and (ii) using a sharp multiscale decomposition; see item (f) below.

Consider a tree graph (or simply tree) θ with k nodes v_1, \dots, v_k and one root \mathbf{r} , which is not considered a node; the tree lines are oriented towards the root (see Fig.1).

The line entering the root is called the root line. We denote by $V(\theta)$ and $\Lambda(\theta)$ the set of nodes and the set of lines in θ , respectively.

- (a) On each node v a label $\boldsymbol{\nu}_v \in \mathbb{Z}^2$, called the *mode label*, is appended.
- (b) To each line ℓ a pair of labels $\eta = (\gamma', \gamma)$ is attached. γ' and γ are called the *left or right component labels*, respectively: $\gamma' \in (1, \dots, 2 + s)$ is associated with the left endpoint of ℓ and $\gamma \in (1, \dots, 2 + s)$ with the right endpoint (in the orientation toward the root, see Fig.1). The label γ' associated with the root line will be denoted by $\gamma(\theta)$.
- (c) Each node v will have $p_v \geq 0$ *entering lines* $\ell_1, \dots, \ell_{p_v}$. Hence with the node v we can associate the left component labels $\gamma'_{v1}, \dots, \gamma'_{vp_v}$ of the entering lines, and an extra label $\gamma_{v0} = \gamma_\ell$ attached to the right endpoint of the line exiting from v . Thus a tensor $\partial_{\gamma_{v0}\gamma_{v1}\dots\gamma_{vp_v}} f_{\boldsymbol{\nu}_v}(\boldsymbol{\beta}_0)$ can be associated with each node v , with ∂_γ denoting the derivative with respect to β_γ if $\gamma > 2$ and multiplication by $i\nu_\gamma$ if $\gamma \leq 2$.
- (d) A *momentum* $\boldsymbol{\nu}_\ell$ is associated with each line $\ell = v'v$ oriented from v to v' : this is a vector in \mathbb{Z}^r defined as $\boldsymbol{\nu}_\ell = \sum_{w \leq v} \boldsymbol{\nu}_w$. The *root momentum*, that is the momentum through the root line, will be denoted by $\boldsymbol{\nu}(\theta)$.
- (e) A *number label* $k_\ell \in \{1, \dots, |\Lambda(\theta)|\}$ is associated with each line ℓ , with $\cup_{\ell \in \Lambda(\theta)} \{k_\ell\} = \{1, \dots, |\Lambda(\theta)|\}$. The number label is used for combinatorial purposes: two trees differing only because of the number labels are still considered distinct.
- (f) Each line ℓ also carries a *scale label* $n_\ell = -1, 0, 1, 2, \dots$: this is a number which determines the size of the *small divisor* $\boldsymbol{\omega} \cdot \boldsymbol{\nu}_\ell$, in terms of an exponentially decreasing sequence $\{\gamma_p\}_{p=0}^\infty$ of positive numbers that we shall introduce in a moment. If $\boldsymbol{\nu}_\ell = \mathbf{0}$ then $n_\ell = -1$. If $|\boldsymbol{\omega} \cdot \boldsymbol{\nu}_\ell| \geq \gamma_0$ then $n_\ell = 0$, and we say that the line ℓ (or else $\boldsymbol{\omega} \cdot \boldsymbol{\nu}_\ell$) is on scale 0. If $\gamma_p \leq |\boldsymbol{\omega} \cdot \boldsymbol{\nu}_\ell| < \gamma_{p-1}$ for some p then $n_\ell = p$, and we say that the line ℓ (or else $\boldsymbol{\omega} \cdot \boldsymbol{\nu}_\ell$) is on scale n . The sequence $\{\gamma_p\}_{p=0}^\infty$ is such that $\gamma_p \in C_0[2^{-p-2}, 2^{-p-1})$ for all $p \geq 0$ and, furthermore, $\boldsymbol{\omega} \cdot \boldsymbol{\nu}$ not only stays bounded below by $C_0|\boldsymbol{\nu}|^{-1}$ (because of the Diophantine condition) but it stays also “far” from the values γ_p for $\boldsymbol{\nu}$ not too large, i.e. for $|\boldsymbol{\nu}|$ at most of order 2^p ; cf. [GG] for a proof of the existence of the sequence (without further assumptions on $\boldsymbol{\omega}$). Precisely,

$$\begin{aligned}
(1) \quad & |\boldsymbol{\omega} \cdot \boldsymbol{\nu}| \geq C_0|\boldsymbol{\nu}|^{-1}, & \mathbf{0} \neq \boldsymbol{\nu} \in \mathbb{Z}^r, \\
(2) \quad & \min_{0 \leq p \leq n} ||\boldsymbol{\omega} \cdot \boldsymbol{\nu}| - \gamma_p| > C_0 2^{-n} & \text{if } n \geq 0, 0 < |\boldsymbol{\nu}| \leq 2^{(n-3)}.
\end{aligned} \tag{2.3}$$

Note that the definition of scale of a line depends on the arbitrary choice of the sequence $\{\gamma_p\}$: we could as well used a sequence scaling as $\gamma_p \sim \gamma^{-p}$, with γ any number > 1 instead of $\gamma = 2$; or we could have used a smooth cutoff function (as in [GG2]) replacing the sharp cutoff function $\mathbb{1}(\gamma_p \leq |\boldsymbol{\omega} \cdot \boldsymbol{\nu}| < \gamma_{p-1})$ implied in the definition above.

(g) The scale labels allow us to define hierarchically ordered clusters. A cluster T of scale n is a maximal connected set of lines ℓ on scale n_ℓ , with $n_\ell \leq n$, containing at least one line on scale n . The lines which are connected to a line of T but do not belong to T are called the *external lines*

of T : according to their orientations, one of them will be called the *exiting line* of T , while all the others will be the *entering lines* of T . All the external lines ℓ are on scales n_ℓ with $n_\ell > n$. The set of lines of T , called the *internal lines* of T , will be denoted by $\Lambda(T)$ and the set of nodes of T will be denoted by $V(T)$.

(h) Not all arrangements of the labels are permitted. The “allowed trees” will have no nodes with $\mathbf{0}$ momentum and with only one entering line and the exiting line also carrying $\mathbf{0}$ momentum. We also discard trees which contain clusters with only one entering line and one exiting line with equal momentum and with no line with $\mathbf{0}$ momentum on the path joining the entering and exiting lines (“self-energy” clusters or “resonances”).

Remark. One can verify that chains of self-energy clusters can actually appear in the initial formal Lindstedt series. One of the main points in [GG1] and [GG2] is to show that if one modifies the series by discarding all chains of self-energy clusters, then the resulting series is convergent (a form of Bryuno’s lemma that appears in KAM theory). In both [GG1] and [GG2] it is shown that, in order to deal with chains of self-energy diagrams one can iteratively resum them into the propagators (i.e. the factors associated with the tree lines in the value of a tree, see below for a definition), that will then turn out to be different from those appearing in the naive formal Lindstedt series (which are simply $(\boldsymbol{\omega} \cdot \boldsymbol{\nu})^{-2}$). Such resummation is the analogue of Dyson’s equations in quantum field theory and the iteratively modified propagator has been, therefore, called the *dressed propagator*. Here there is further freedom in the choice of the self-energy clusters. The idea is that the self-energy clusters must include the “diverging contributions” affecting the initial formal power series. But if we change the definition of self-energy clusters by adding to the class a new class of non-diverging clusters, the construction can be shown to go through as well. We find convenient the specific choice above but this is of course not necessary. This is the second arbitrary choice we do in the iterative construction of the resummed series. It can fuel doubts about the uniqueness of the result which can only be dismissed by further arguments (like the Borel summability that we are proving).

The set of all allowed trees with labels $\gamma(\theta) = \gamma$ and $\boldsymbol{\nu}(\theta) = \boldsymbol{\nu}$ is denoted by $\Theta_{\boldsymbol{\nu}\gamma}$ (this is the set appearing in (2.2)). The labels described above are used to define the *value* $\text{Val}(\theta)$ of a (decorated) tree $\theta \in \Theta_{\boldsymbol{\nu}\gamma}$: this is a number obtained by multiplying the following factors:

- (1) a factor $F_v = \partial_{\gamma_{v0}\gamma'_{v1}\dots\gamma'_{vp_v}} f_{\boldsymbol{\nu}_v}(\boldsymbol{\beta}_0)$, called the *node factor*, per each node v ;
- (2) a factor $g_\ell^{[n_\ell]} = g_{\gamma'_\ell, \gamma_\ell}^{[n_\ell]}(\boldsymbol{\omega} \cdot \boldsymbol{\nu}_\ell; \eta)$, called the *propagator*, per each line of scale n_ℓ , momentum $\boldsymbol{\nu}_\ell$ and component labels $\gamma'_\ell, \gamma_\ell$, see items (I)–(V) below for a definition.

The value is then defined as

$$3.2 \quad \text{Val}(\theta) = \frac{1}{k!} \left(\prod_{v \in V(\theta)} F_v \right) \left(\prod_{\ell \in \Lambda(\theta)} g_\ell^{[n_\ell]} \right), \quad (2.4)$$

where it should be noted that all labels γ (of the tensors F_v and of the matrices $g_\ell^{[n_\ell]}$) appear repeated twice because they appear in the propagators as well as in the tensors associated with the nodes, with the exception of the label γ associated with the left endpoint of the line ending in the root (as the root is not a node and therefore there is no tensor associated with it).

Adopting the convention of summation over repeated component labels $\text{Val}(\theta)$ depends on the root label $\gamma(\theta) = \gamma = 1, \dots, 2 + s$ so that it defines a vector in \mathbb{C}^{2+s} .

The recursive definition of the propagators is such that the series in (2.2) is convergent and gives the ν -th Fourier component of the function $\mathbf{h}(\psi, \eta)$ in Sect. 1. The definition of propagators we adopt here is the same introduced in [GG2]: the definition in [GG1] is slightly different (see Appendix A2), but it has the drawback that it is specific for hyperbolic resonances, while the definition in [GG2] can be (expected to be) extended also to the theory of elliptic resonances and, therefore, might turn out to be useful in view of possible extensions of the main results of this work to elliptic resonances.

(I) For $n = -1$ the propagator of the line ℓ is defined as the block matrix

$$3.4 \quad g_\ell^{[-1]} \stackrel{def}{=} \begin{pmatrix} 0 & 0 \\ 0 & (-\partial_{\beta}^2 f_0(\beta_0))^{-1} \end{pmatrix}. \quad (2.5)$$

(II) For $n = 0$, if the line ℓ carries a momentum ν and if $x \stackrel{def}{=} \omega \cdot \nu$, the propagator is the matrix

$$3.5 \quad g_\ell^{[0]} = g^{[0]}(x; \eta) = \frac{\eta^2}{x^2 + \eta^2 M_0}, \quad (2.6)$$

with $M_0 \stackrel{def}{=} \begin{pmatrix} 0 & 0 \\ 0 & -\partial_{\beta}^2 f_0(\beta_0) \end{pmatrix}$. By the assumptions of Proposition 2 one has $M_0 \geq 0$.

(III) For $n > 0$ the propagator is the matrix

$$3.6 \quad g_\ell^{[n]} = g^{[n]}(x; \eta) = \frac{\eta^2}{x^2 + \mathcal{M}^{[\leq n]}(x; \eta)}. \quad (2.7)$$

with $\mathcal{M}^{[\leq n]}(x; \eta) = \mathcal{M}^{[0]}(x; \eta) + \mathcal{M}^{[1]}(x; \eta) + \dots + \mathcal{M}^{[n]}(x; \eta)$, where $\mathcal{M}^{[0]}(x; \eta) = \eta^2 M_0$, whereas $\mathcal{M}^{[j]}(x; \eta)$, $j \geq 1$, are matrices, called *self-energy matrices*, whose expansion in η starts at order η^4 and are defined as described in the next two items.

(IV) Let T be a self-energy cluster on scale n (see item (h) above) and let us define the matrix¹ $\mathcal{V}_T(\omega \cdot \nu; \eta)$ as

$$3.7 \quad \mathcal{V}_T(\omega \cdot \nu; \eta) = -\frac{\eta^2}{|\Lambda(T)|!} \left(\prod_{v \in V(T)} F_v \right) \left(\prod_{\ell \in \Lambda(T)} g_\ell^{[n_\ell]} \right), \quad (2.8)$$

where, necessarily, $n_\ell \leq n$ for all $\ell \in \Lambda(T)$. The matrix (2.8) will be called the *self-energy value* of T . The set of the self-energy clusters with value proportional to η^{2k} , hence with $k - 1$ internal lines with $n_\ell \geq 0$, and with maximum scale label n will be denoted $\mathcal{S}_{k,n}^{\mathcal{R}}$.

(V) The *self-energy matrices* $\mathcal{M}^{[n]}(x; \eta)$, $n \geq 1$, are defined recursively for $|x| \leq \gamma_{n-1}$ (i.e. for x on scale $\geq n$) as

$$3.8 \quad \mathcal{M}^{[n]}(x; \eta) = \sum_{k=2}^{\infty} \sum_{T \in \mathcal{S}_{k,n-1}^{\mathcal{R}}} \mathcal{V}_T(x; \eta), \quad (2.9)$$

where the self-energy values are evaluated by means of the propagators on scales p , with $p = -1, 0, 1, \dots, n - 1$.

Remarks.

(1) With respect to [GG2] the second argument of the propagators (and of the self-energy values

¹ This is a matrix because the self-energy cluster inherits the labels γ, γ' attached to the left of the entering line and to the right of the exiting line.

and matrices) has been denoted η instead of ε ; we recall that the variable η^2 appearing here is the same as the variable ε appearing in [GG1] and [GG2]. We make this choice because it is natural to study Borel summability in η and not in $\varepsilon = \eta^2$.

(2) The association of the factors η^2 with the lines themselves rather than with the nodes (as in [GG1] and [GG2]) will be more convenient when considering the Borel transforms of the involved quantities.

(3) The multiscale decomposition used in [GG2] may look quite different from the one we are using here, but this is not really so. First, even though the decomposition in [GG2] was based on the propagator divisors $\Delta^{[n]}(x; \varepsilon) = \min_j |x^2 - \lambda_j^{[n]}(\varepsilon)|$, where the self-energies $\lambda_j^{[n]}(\varepsilon)$ were defined recursively in terms of the self-energy matrices, in the case of hyperbolic tori one has identically $\Delta^{[n]}(x; \varepsilon) = x^2$: indeed all self-energies which are non-zero are strictly negative. Then the only real difference is that here we are using a sharp decomposition instead of a smooth one, but the latter is not a relevant difference. In fact the choice of the sequence $\{\gamma_p\}_{p=0}^\infty$ implies that the lines appearing in the groups of graphs that will be collected together to exhibit the necessary cancellations have currents ν such that $\nu \cdot \omega$ stays relatively far from the extremes of the intervals $[\gamma_{p+1}, \gamma_p]$ that define the scale labels, and this allows us to use a sharp multiscale decompositions instead of the smooth one used in [GG2]. In other words this change with respect to [GG2] is done only to avoid introducing partitions of unity by smooth functions and the related discussions.

Therefore the expression (2.2) makes sense and in fact the function \mathbf{h} mentioned in Proposition 1 is exactly the Fourier sum of the r.h.s. of (2.2). In particular in [GG2] it was proved that the Fourier sum of the r.h.s. of (2.2) satisfies the properties (i)–(iii) in Proposition 1.

sec.4

3. Integral representation of the resummed Lindstedt series

Given the definitions of Sect. 2 consider the function $\mathbf{h}^{(N)}$ defined in the same way as \mathbf{h} but restricting the sum in (2.2) to the trees containing only lines of scale $n \leq N$.

The functions $\mathbf{h}^{(N)}$ have the “same” convergence and analyticity properties of the functions \mathbf{h} and the same bounds on the Taylor coefficients at the origin. Moreover $\mathbf{h}^{(N)} \xrightarrow{N \rightarrow \infty} \mathbf{h}$: this is a consequence of the intermediate steps in the proof of the above proposition in [GG2], as the strategy of the proof is to define $\mathbf{h}^{(N)}$ making sure that the convergence and analyticity properties are uniform in N . In fact $\mathbf{h}^{(N)}$ is even analytic in η near the origin for $|\eta| \leq \eta_N$ (but $\eta_N \xrightarrow{N \rightarrow \infty} 0$).

Therefore the functions $\mathbf{h}^{(N)}$ are trivially Borel summable, and have an entire Borel transform, but the growth at $p \rightarrow +\infty$ of their Borel transforms is N -dependent while, to show Borel summability of \mathbf{h} , uniform estimates are needed. This section is devoted to a first attempt at such bounds which uses minimally the informations on the resummed series that can be gathered from [GG2], i.e. the convergence properties just mentioned.

The Borel transform of the functions $\mathbf{h}^{(N)}(\psi, \eta)$ is an entire function that can be written for p real and positive as

$$4.1 \quad (\mathbf{h}^{(N)})_B(\psi, p) = \mathcal{L}^{-1}h(\psi, p) = \int_{\eta_N^{-1}-i\infty}^{\eta_N^{-1}+i\infty} e^{pz} \mathbf{h}^{(N)}\left(\psi, \frac{1}{z}\right) \frac{dz}{2\pi i}, \quad p \in \mathbb{R}^+, \quad (3.1)$$

where η_N is the convergence radius of $\mathbf{h}^{(N)}(\psi, \eta)$. The key remark is that we also know that by property (ii) in Proposition 1 (actually by the same property for $\mathbf{h}^{(N)}$ that follows from the construction in [GG2]) the function $\mathbf{h}^{(N)}(\psi, \frac{1}{z})$ is analytic for $|z| > 2\eta_0^{-1}$, so that the integral in

(3.1) can be shifted to a contour on the vertical line with abscissa $\bar{\rho} > 2\eta_0^{-1}$, i.e. with N -independent abscissa. Therefore

$$4.2 \quad (\mathbf{h}^{(N)})_B(\boldsymbol{\psi}, p) = \int_{\bar{\rho}-i\infty}^{\bar{\rho}+i\infty} e^{pz} \mathbf{h}^{(N)}\left(\boldsymbol{\psi}, \frac{1}{z}\right) \frac{dz}{2\pi i}, \quad p \in \mathbb{R}^+, \quad (3.2)$$

and, for all N , the function $\mathbf{h}^{(N)}(\boldsymbol{\psi}, \frac{1}{z})$ is uniformly bounded by $O(\frac{1}{|z|^2})$ on the integration contour, because, by property (iii) in Proposition 1, \mathbf{h} is twice differentiable at the origin along any path contained in \mathcal{D} (in particular along the circular path $\text{Re } \eta^{-1} = \rho$). Hence the latter boundedness property of $\mathbf{h}^{(N)}(\boldsymbol{\psi}, \frac{1}{z})$ and (3.2) imply the bound, for $p > 0$ and for a suitable constant C ,

$$4.3 \quad |(\mathbf{h}^{(N)})_B(\boldsymbol{\psi}, p)| \leq C e^{\bar{\rho}p}, \quad \forall p \in \mathbb{R}^+, \quad (3.3)$$

and for all $\bar{\rho} > 2\eta_0^{-1}$. The existence of the limit $\lim_{N \rightarrow \infty} \mathbf{h}^{(N)}(\boldsymbol{\psi}, \frac{1}{z}) = \mathbf{h}(\boldsymbol{\psi}, \frac{1}{z})$ for $\frac{1}{|z|}$ small (by [GG2]) implies existence of the limit $\mathbf{F}(\boldsymbol{\psi}, p)$ as $N \rightarrow \infty$ of $(\mathbf{h}^{(N)})_B(\boldsymbol{\psi}, p)$ for $p \in \mathbb{R}^+$ and $\mathbf{F}(\boldsymbol{\psi}, p)$ satisfies the bound (3.3) on \mathbb{R}^+ .

Hence the functions $\mathbf{h}(\boldsymbol{\psi}, \eta)$ can be expressed, for $0 \leq \eta < \eta_0$, as

$$4.4 \quad \mathbf{h}(\boldsymbol{\psi}, \eta) = \int_0^\infty e^{-p/\eta} \left(\lim_{N \rightarrow \infty} (\mathbf{h}^{(N)})_B(\boldsymbol{\psi}, p) \right) dp = \int_0^\infty e^{-p/\eta} \mathbf{F}(\boldsymbol{\psi}, p) dp, \quad (3.4)$$

which provides us with an integral representation of the resummed series and shows that the resummation (2.2) generates a Borel sum of the formal Lindstedt series *provided* $\mathbf{F}(\boldsymbol{\psi}, p)$ can be shown to be analytic in a neighborhood of the positive axis, as required by the very definition of Borel summability, see property (ii) in Definition 1 of Section 1.2.

sec.5

4. Borel summability

By the remark at the end of Sect. 3 Borel summability of \mathbf{h} will be established once the natural candidate for its Borel transform, namely the function \mathbf{F} in (3.4), is shown to be analytic in a region containing the positive real axis. This will be done here assuming, for simplicity, that the perturbation f is a trigonometric polynomial. The general case of an analytic f is slightly more involved and will be treated in Appendix A1.

Assume, inductively, that $\forall x = \boldsymbol{\omega} \cdot \boldsymbol{\nu}$ the functions $(\mathcal{M}^{[\leq k]})_B(x; p)$ are entire functions of p and

$$5.1 \quad \|(\mathcal{M}^{[\leq k]})_B(x; p)\| \leq \Gamma |p| e^{d_k |p|}, \quad \text{for all } x \text{ on scale } k', \text{ with } k' \geq k \geq 0, \quad (4.1)$$

for all p complex. The matrix norm in (4.1) is $\|\mathcal{M}\| = \max_j \sum_i |\mathcal{M}_{ij}|$. Note that $(\mathcal{M}^{[0]})_B(p) = pM_0$ (so that the inductive assumption in (4.1) is valid at the first step $k = 0$ with $d_0 = 0$ if $\Gamma \geq \|M_0\|$) and $(g^{[0]})_B(x; p)$ is entire and it is given by

$$5.2 \quad (g^{[0]})_B(x; p) = \frac{1}{x^2} \frac{\sin p \sqrt{M_0 x^{-2}}}{\sqrt{M_0 x^{-2}}} \Rightarrow \|(g^{[0]})_B(x; p)\| \leq \frac{|p|}{x^2} e^{|p|c_0}, \quad \forall |x| \geq \gamma_0, \quad (4.2)$$

with $c_0 = \sqrt{\|M_0\|}/\gamma_0$.

Supposing the inductive assumption to be valid for $k = 1, \dots, N-1$ we remark that this implies a bound on $(g^{[k]})_B(x; p)$ for $k = 1, \dots, N-1$ via the expansion

$$5.3 \quad (g^{[k]})_B(x; p) = \left(\frac{\eta^2}{x^2} \sum_{m=0}^{\infty} \left(\frac{-\mathcal{M}^{[\leq k]}(x; \cdot)}{x^2} \right)^m \right)_B. \quad (4.3)$$

Taking the Borel transform and performing all convolutions along a straight line from 0 to p , for x on scale $1 \leq k \leq N-1$, we get

$$\begin{aligned} \|(g^{[k]})_B(x; p)\| &\leq \frac{1}{x^2} \sum_{m=0}^{\infty} |p| * \frac{(\Gamma|p|e^{d_k|p|})^{*m}}{x^{2m}} \leq \frac{|p|}{x^2} \sum_{m=0}^{\infty} e^{d_k|p|} \frac{\Gamma^m |p|^{2m}}{(2m)!} \frac{1}{x^{2m}} \leq, \\ &\leq \frac{|p|}{x^2} e^{d_k|p| + \Gamma^{1/2} \gamma_k^{-1} |p|} \equiv \frac{|p|}{x^2} e^{c_k|p|} \end{aligned} \quad (4.4)$$

with $f^{*m} \stackrel{def}{=} f * \dots * f$ m times and $c_k = d_k + \Gamma^{1/2} \gamma_k^{-1}$.

Then $(\mathcal{M}^{[\leq N]} - \mathcal{M}^{[0]})_B(x; p)$ is estimated via (2.9), that is

$$\begin{aligned} &\|(\mathcal{M}^{[\leq N]} - \mathcal{M}^{[0]})_B(x; p)\| \leq \\ &\leq \sum_{k=2}^{\infty} \sum_{T \in \cup_{j=0}^{N-1} \mathcal{S}_{k,j}^{\mathcal{R}}} \frac{1}{|\Lambda(T)|!} |p| * \left(\prod_{\substack{\ell \in \Lambda(T) \\ n_{\ell} \geq 0}}^* \frac{1}{x_{\ell}^2} |p| e^{c_{n_{\ell}} |p|} \right) \left(\prod_{\substack{\ell \in \Lambda(T) \\ n_{\ell} = -1}} \|g_{\ell}^{[-1]}\| \right) \left(\prod_{v \in V(T)} \|F_v\| \right), \end{aligned} \quad (4.5)$$

where $x_{\ell} = \omega \cdot \nu_{\ell}$, the \prod^* is a convolution product and

$$\|F_v\| = \max_{\gamma'_{v1}, \gamma'_{v2}, \dots, \gamma'_{vpv}} \sum_{\gamma_{v0}} |(F_v)_{\gamma_{v0}, \gamma'_{v1}, \gamma'_{v2}, \dots, \gamma'_{vpv}}|. \quad (4.6)$$

Hence bounding $|p| * \left(\prod^* |p| e^{c_{n_{\ell}} |p|} \right)$ by $e^{c_{N-1} |p|} |p|^{2k-1} / (2k-1)!$ and using the estimates in [GG2] to control the sum over the self-energy clusters, the bound becomes

$$\|(\mathcal{M}^{[\leq N]})_B(x; p)\| \leq \Gamma |p| + \sum_{k=2}^{\infty} \Gamma^{2k} \frac{|p|^{2k-1}}{(2k-1)!} e^{c_{N-1} |p|} \leq \Gamma |p| e^{d_N |p|}, \quad (4.7)$$

where Γ is a suitable constant derived in [GG1] and $d_N = c_{N-1} + \Gamma$.

Remark. The step leading from (4.5) to (4.7) is non-trivial and the possibility of bounding the small divisors x_{ℓ}^2 and the sum over self-energy clusters, after defining the propagators as above, is the main technical aspect of the work in [GG2]. We take the existence of Γ from [GG2]. We do not repeat here the analysis performed in Sects. 5 and 6 (and the corresponding Appendices) of [GG2]: the constant Γ^2 has been called $\bar{\varepsilon}^{-1}$ in Theorem 1 of [GG2].

Thus the inductive assumption holds for all N , the constants c_N, d_N can be taken $c2^N$ for some c , and for all x on scale N one has

$$\|(g^{[N]})_B(x; p)\| \leq \frac{|p|}{x^2} e^{2^N c |p|}. \quad (4.8)$$

This leads to a bound on $(\mathbf{h}^{(N)})_B(\psi, p)$, via (2.4) and (2.2):

$$\begin{aligned} |(\mathbf{h}^{(N)})_B(\psi, p)| &\leq \sum_{k=1}^{\infty} \sum_{\nu} \sum_{\Theta_{\nu, \gamma}} \frac{1}{|\Lambda(\theta)|!} \left(\prod_{\substack{\ell \in \Lambda(\theta) \\ n_{\ell} \geq 0}}^* \frac{1}{x_{\ell}^2} |p| e^{c_{n_{\ell}} |p|} \right) \left(\prod_{\substack{\ell \in \Lambda(T) \\ n_{\ell} = -1}} \|g_{\ell}^{[-1]}\| \right) \left(\prod_{v \in V(T)} \|F_v\| \right) \leq \\ &\leq \sum_{k=1}^{\infty} \Gamma^{2k} \frac{|p|^{2k-1}}{(2k-1)!} e^{c^{2^k} |p|} \leq \sum_{k=1}^{\infty} \Gamma^{2k} \frac{|p|^{2k-1}}{(2k-1)!} e^{c' k |p|} \leq \Gamma^2 |p| e^{\Gamma |p| e^{c' |p|}}, \end{aligned} \quad (4.9)$$

because the maximum scale n_{\max} of the lines of a graph with k lines can be at most $\log_2(bk)$ by our assumption that f is a trigonometric polynomial: in fact the maximum momentum on a line can be $|\nu| \leq b_0 k$ (for f a trigonometric polynomial), so that the smallest x can be $\frac{C_0}{b_0 k}$ if C_0 and $\tau = 1$ are the Diophantine constants, hence the scale of x can be at most $\log_2(b_0 k/4)$ and $c2^{n_{\max}} \leq c'k$ for a suitable c' .

Therefore the functions $(\mathbf{h}^{(N)})_B(\boldsymbol{\psi}; p)$ are entire and bounded by

$$5.9 \quad |(\mathbf{h}^{(N)})_B(\boldsymbol{\psi}; p)| \leq \Gamma^2 |p| e^{\Gamma |p| e^{c' |p|}} \quad (4.10)$$

independently of N . Since for $|p|$ small the functions $(\mathbf{h}^{(N)})_B(\boldsymbol{\psi}; p)$ converge to $\mathbf{F}(p)$, simply because of the bound $|\frac{1}{k!} \partial^k \mathbf{h}(\boldsymbol{\psi}, 0)| \leq DC^k k!$ on the Taylor coefficients at the origin of $\mathbf{h}(\boldsymbol{\psi}, \eta)$, it follows that $\mathbf{F}(\boldsymbol{\psi}; p)$ is entire in p and everywhere the limit of $(\mathbf{h}^{(N)})_B(\boldsymbol{\psi}; p)$. Therefore (by Vitali's theorem) $\mathbf{F}(\boldsymbol{\psi}; p)$ is not only holomorphic in p near 0 (which is a result on the Lindstedt series at the origin) but it is entire. In particular it is holomorphic around the real axis and therefore by (3.4) $\mathbf{h}(\boldsymbol{\psi}; \eta)$ is Borel summable.

sec.6

5. Concluding remarks

(1) It is interesting to stress that the recursive bounds on the Borel transform of the propagators in (4.4) have been derived without making use of the cancellations that played such an essential role in the theory in [GG2] (and [GG1]) and by “undoing” at each step the resummations which led to the construction of \mathbf{h} and to the proof of Proposition 1, see (4.3) above. However the properties of \mathbf{h} and the result of Proposition 1 have played a key role in the derivation of (3.3) and (3.4). Without the uniform bounds (in N) on the convergence radii in p of the series expressing $(\mathbf{h}^{(N)})_B$, which depend on the cancellations and on the resummations, the bounds in Sect. 4 or, in the non-trigonometric case, of Appendix A1, would remain the same but they would be useless for our purposes of establishing (3.4) and the Borel summability.

(2) The assumption that f is a trigonometric polynomial has been heavily exploited and an extra idea is necessary to deal with the more general case of analytic f : this is discussed in Appendix A1.

(3) It has been remarked above that the resummation procedure is based on several arbitrary *a priori* choices which therefore may lead to the existence of several solutions of \mathbf{h} with the same asymptotic series at $\varepsilon = 0$. All the choices in [GG2], as well as that used in [GG1] (see Appendix A2), lead however to a Borel summable series: this proves that all solutions coincide and the results of the resummations are independent of the particular choices *provided* the Diophantine constant τ is $\tau = 1$ (hence $r = 2$ and $\boldsymbol{\omega}$ is a Diophantine vector with $\tau = 1$, e.g. ω_1/ω_2 is a quadratic irrational).

(4) If $\tau > 1$, hence if $r > 2$, the problem of Borel summability and of independence of the result from the summation method remains open.

(5) The existence or non-existence of solutions \mathbf{h} which are C^∞ at the origin and give solutions to the equations of motion but which are not Borel summable is also an open problem. Note that *even in the case of non-resonant motions* the problem of the uniqueness at fixed $\boldsymbol{\omega}$ is not trivial, and also the recent results in [BT] do not exclude the possibility of other quasi-periodic motions besides those constructed through the KAM algorithm.

(6) The case $\varepsilon < 0$ with β_0 a minimum point for $f_0(\boldsymbol{\beta})$, i.e. of elliptic motions is quite different. We have used on purpose the resummation technique of [GG2] which works for hyperbolic as well

as for elliptic resonances, but the present results still only apply to the hyperbolic case and it is not clear whether the above techniques can be extended to prove Borel summability of the parametric equations for \mathbf{h} in elliptic cases.

app.A1

Appendix A1. Analytic, non-polynomial perturbations

In this Appendix we want to prove real analyticity of $\mathbf{F}(\boldsymbol{\psi}, p)$ in the case f is a generic analytic function (rather than a trigonometric polynomial); in this case the function $\mathbf{F}(\boldsymbol{\psi}, p)$ will turn out not be entire but only analytic in a strip of width $2\kappa > 0$ around the real axis.

The proof of this claim will be based on an inductive assumption on $(g^{[n]})_B(x; p)$ formulated by introducing the matrix $\tilde{g}^{[n]}(x; \eta) = \frac{\eta^2}{x^2 + \mathcal{M}^{[\leq n]}(x; \eta)}$ for all $|x| < \gamma_{n-1}$. Note that if $\chi_n(x) \stackrel{\text{def}}{=} \mathbb{1}(\gamma_n \leq |x| < \gamma_{n-1})$ is the indicator of the scale of x , the propagator $g^{[n]}(x; \eta)$ is given by $g^{[n]}(x; \eta) = \chi_n(x) \tilde{g}^{[n]}(x; \eta)$. Furthermore the matrices $g^{[n]}(x; \eta)$ satisfy the recursive equations

$$A1.2 \quad (\tilde{g}^{[n]}(x; \eta))^{-1} = (\tilde{g}^{[n-1]}(x; \eta))^{-1} + \eta^{-2} \mathcal{M}^{[n]}(x; \eta), \quad \forall |x| < \gamma_{n-1}. \quad (A1.1)$$

We suppose, inductively, that for $\kappa_0 = \sqrt{\|\overline{M_0}\|}$ one has

$$A1.3 \quad \|(\tilde{g}^{[n]})_B(x; p)\| \leq K_0 \frac{|p|}{x^2} e^{(c_n + c'_n |x|^{-1/2})|p| + \kappa_0 |\text{Im } p| |x|^{-1}}, \quad \forall |x| < \gamma_{n-1}, \quad n \geq 0. \quad (A1.2)$$

Note that for $n = 0$ the explicit expression of $(\tilde{g}^{[0]})_B$ given by (4.2) implies that (A1.2) is valid with $K_0 = \sqrt{s}$ (where s is the dimension of the non trivial block in M_0) and $c_0 = c'_0 = 0$. The constant K_0 comes from our choice of the matrix norm $\|M\| = \max_j \sum_i |M_{ij}|$: with this choice for any $d \times d$ matrix we have $d^{-1/2} \|M\|_2 \leq \|M\| \leq d^{1/2} \|M\|_2$ where $\|\cdot\|_2$ is the spectral norm, so that in particular $\|\sin M/M\|, \|\cos M\| \leq d^{1/2}$.

Assuming (A1.2) for $n \leq N-1$ and performing all convolutions along the straight line from the origin to p , we get the following bound on $(\mathcal{M}^{[N]})_B(x; p)$:

$$A1.4 \quad \begin{aligned} & \|(\eta^{-2} \mathcal{M}^{[N]})_B(x; p)\| \leq \\ & \leq \sum_{k=2}^{\infty} \sum_{T \in \mathcal{S}_{k, N-1}^{\mathcal{R}}} \frac{1}{|\Lambda(T)|!} \left(\prod_{\substack{\ell \in \Lambda(T) \\ n_\ell \geq 0}}^* K_0 \frac{|p|}{x^2} e^{(c_{n_\ell} + c'_{n_\ell} \gamma_{n_\ell}^{-1/2})|p| + \kappa_0 |\text{Im } p| \gamma_{n_\ell}^{-1}} \right) \\ & \quad \left(\prod_{\substack{\ell \in \Lambda(T) \\ n_\ell = -1}} \|g_\ell^{[-1]}\| \right) \left(\prod_{v \in V(T)} \|F_v\| \right) \\ & \leq \sum_{k=2}^{\infty} \Gamma^{2k} \frac{|p|^{2k-3}}{(2k-3)!} e^{(c_{N-1} + c'_{N-1} \gamma_{N-1}^{-1/2})|p| + \kappa_0 |\text{Im } p| \gamma_{N-1}^{-1}} e^{-2\kappa_0 2^N} \leq D_0 |p| e^{d_N |p|} e^{-\kappa_0 2^N}, \end{aligned} \quad (A1.3)$$

where Γ and κ are deducible from Appendix A3 of [GG2], $D_0 = \Gamma^3$, $d_N = c_{N-1} + c'_{N-1} \gamma_{N-1}^{-1/2}$ and, using that $\gamma_{N-1}^{-1} \leq 4C_0^{-1} 2^N$, we chose $|\text{Im } p|$ so small that $4\kappa_0 C_0^{-1} |\text{Im } p| \leq \kappa$.

Using (A1.3) and (A1.1) we can get a bound on $(\tilde{g}^{[N]})_B(x; p)$:

$$A1.5 \quad \begin{aligned} & \|(\tilde{g}_N)_B(x; p)\| \leq \left(K_0 \frac{|p|}{x^2} e^{(c_{N-1} + c'_{N-1} |x|^{-1/2})|p| + \kappa_0 |\text{Im } p| |x|^{-1}} \right)^* \\ & * \sum_{k=0}^{\infty} \left[\left(K_0 \frac{|p|}{x^2} e^{(c_{N-1} + c'_{N-1} |x|^{-1/2})|p| + \kappa_0 |\text{Im } p| |x|^{-1}} \right)^* \left(D_0 |p| e^{d_N |p|} e^{-\kappa_0 2^N} \right)^* \right]^k, \end{aligned} \quad (A1.4)$$

and, since $|p| * (|p| * |p|)^{*k} = \frac{|p|^{4k+1}}{(4k+1)!}$ for $k \geq 0$, the k -th term in the sum is bounded by

$$A1.6 \quad K_0 \frac{(K_0 D_0 e^{-\kappa 2^N})^k}{(4k+1)!} \frac{|p|}{x^2} \frac{|p|^{4k}}{x^{2k}} e^{(d_N + c'_{N-1}|x|^{-1/2})|p| + \kappa_0 |\operatorname{Im} p| |x|^{-1}}. \quad (A1.5)$$

Summing (A1.5) over $k \geq 0$ and comparing the result with the inductive assumption (A1.2), we see that we can take $c_N = d_N = c_{N-1} + c'_{N-1} \gamma_{N-1}^{-\alpha}$ and $c'_N = c'_{N-1} + K_0 D_0 e^{-\kappa 2^N}$. Solving the iterative equations for c_N, c'_N , we see that, for x on scale n , $(\tilde{g}^{[n]})_B(x; p)$ can be bounded as

$$A1.6a \quad \|(\tilde{g}^{[n]})_B(x; p)\| \leq K_0 \frac{|p|}{x^2} e^{c 2^{n/2} |p| + \kappa_1 |\operatorname{Im} p| 2^n}, \quad \gamma_n \leq |x| < \gamma_{n-1}, \quad n \geq 0, \quad (A1.6)$$

for some constants c, κ_1 .

Plugging this bound into the expansion for $\mathbf{h}_B^{(N)}(\psi, p)$, denoting by $N(\theta)$ the maximal scale in θ and choosing $|\operatorname{Im} p|$ small enough, we finally get

$$A1.7 \quad \begin{aligned} & |(\mathbf{h}^{(N)})_B(\psi, p)| \leq \\ & \leq \sum_{k=1}^{\infty} \sum_{\nu} \sum_{n_0 \geq 0} \sum_{\substack{\theta \in \Theta_{\nu, \gamma} \\ N(\theta) = n_0}} \frac{1}{|\Lambda(\theta)|!} \left(\prod_{\substack{\ell \in \Lambda(\theta) \\ n_{\ell} \geq 0}}^* \frac{K_0}{x_{\ell}^2} |p| e^{c 2^{n_{\ell}/2} |p| + \kappa_1 |\operatorname{Im} p| 2^{n_{\ell}}} \right) \\ & \left(\prod_{\substack{\ell \in \Lambda(T) \\ n_{\ell} = -1}} \|g_{\ell}^{[-1]}\| \right) \left(\prod_{v \in V(T)} \|F_v\| \right) \leq \\ & \leq \sum_{k=1}^{\infty} \sum_{n_0 \geq 0} \Gamma^{2k} \frac{|p|^{2k-1}}{(2k-1)!} e^{c 2^{n_0/2} |p| + \kappa_1 |\operatorname{Im} p| 2^{n_0}} e^{-2\kappa 2^{n_0}} \leq \Gamma' |p| e^{c' |p|^2}, \end{aligned} \quad (A1.7)$$

for some new constants Γ', c' . Performing the limit $N \rightarrow \infty$ in (A1.7), we see that $\mathbf{F}(\psi, p)$ satisfies the same bound

$$A1.8 \quad |\mathbf{F}(\psi, p)| \leq \Gamma' |p| e^{c' |p|^2}, \quad \text{for } |\operatorname{Im} p| \leq \sigma, \quad (A1.8)$$

for some σ small enough, so that $\mathbf{F}(\psi, p)$ is analytic (in p) in a strip and therefore $\mathbf{h}(\psi, \eta)$ is Borel summable (in η).

app.A2

Appendix A2. Comparison with the method of [GG1]

In this Appendix we briefly discuss how the function \mathbf{h} constructed in [GG1] can be identified with the Borel summable function of Proposition 1. By the uniqueness in the class of Borel summable functions, it is enough to prove that also the function \mathbf{h} of [GG1] is Borel summable.

We begin by reviewing the differences of the construction envisaged in [GG1] with respect to that of [GG2]. Trees, labels and clusters are defined in the same way as in items (a) to (h) of Section 2. What changes is the definition of the propagators, which is iterative. By writing $x = \omega \cdot \nu_{\ell}$, $\nu_{\ell} \neq \mathbf{0}$, we set $g_{\ell}^{[0]} = 1/x^2$, and, for $k \geq 1$,

$$A2.1 \quad g_{\ell}^{[k]} = \frac{\eta^2}{x^2 + M^{[k]}(x; \eta)}. \quad (A2.1)$$

with $M^{[k]}(x; \eta)$ defined as

$$A2.2 \quad M^{[k]}(x; \eta) = \sum_T \mathcal{V}_T(x; \eta), \quad (A2.2)$$

where the sum is restricted to the self-energy clusters T with scale $n_T \geq n + 3$, where n is such that $\gamma_{n-1} \leq |x| < \gamma_n$, and the self-energy value is given by

$$A2.3 \quad \mathcal{V}_T(\boldsymbol{\omega} \cdot \boldsymbol{\nu}; \eta) = -\frac{\eta^2}{|\Lambda(T)|!} \left(\prod_{v \in V(T)} F_v \right) \left(\prod_{\ell \in \Lambda(T)} g_\ell^{[k-1]} \right). \quad (A2.3)$$

Note that k labels the iterative step and, in principle, it has no relation with the scale n of x . However the matrices $M^{[k]}(x; \eta)$ are obtained from resummations of self-energy clusters with height up to $k = n$ (for definitions and details we refer to [GG1], where the self-energy clusters are called self-energy graphs). Hence they stop flowing at $k = n$ if x is on scale n : this means that $M^{[k]}(x; \eta) = M^{[n]}(x; \eta)$ as soon as $k \geq n$.

Then $\mathbf{h}^{(N)}$ will be expressed in terms of trees as in (2.2), if $\text{Val}(\theta)$ is defined as in (2.4), with $g_\ell^{[n_\ell]}$ replaced with $g_\ell^{[N]}$, and $\mathbf{h}^{(N)}$ is obtained as the limit of $\mathbf{h}^{(N)}$ as $N \rightarrow \infty$.

Let us consider for simplicity the case of polynomial perturbations. Then we can proceed as in Sec. 4, and prove by induction the bound

$$A2.4 \quad \|(M^{[k]})_B(x; p)\| \leq \Gamma |p| e^{d_k |p|}, \quad \text{for all } x, \quad (A2.4)$$

for all p complex. Supposing inductively the bound (A2.4) we obtain that the Borel transform of $g_\ell^{[k]}$ can be bounded as

$$A2.5 \quad \|(g^{[k]})_B(x; p)\| \leq \frac{|p|}{x^2} e^{c_k |p|}. \quad (A2.5)$$

This is trivial for $k = 0$, as $(g^{[0]})_B(x; p) = p/x^2$, while it follows from (A2.4) for $n \geq 1$ by the inductive hypothesis (and it can be proved as the analogous bound in Sec. 4). Therefore we can write $M^{[N]}(x; \eta)$ according to (A2.2), and its Borel transform $(M^{[N]})_B(x; p)$ can be computed and bounded as done in (4.5) and (4.8), so that at the end the bound (A2.4) is obtained for $k = N$. In particular the bounds (A2.5) on the propagators imply that $\mathbf{h}^{(N)}$ admits the same bound (4.10) found in Sec. 4. Therefore we can take the limit for $N \rightarrow \infty$, and Borel summability for \mathbf{h} follows.

References

- 1[BT] H. Broer, F. Takens, *Unicity of KAM tori*, Preprint, Groningen, 2005.
- 2[GBG] G. Gallavotti, F. Bonetto, G. Gentile, *Aspects of ergodic, qualitative and statistical theory of motion*, Texts and Monographs in Physics, Springer, Berlin, 2004.
- 3[GG] G. Gallavotti, G. Gentile, *Majorant series convergence for twistless KAM tori*, Ergodic Theory and Dynamical Systems **15** (1995), 857–869.
- 4[GG1] G. Gallavotti, G. Gentile, *Hyperbolic low-dimensional invariant tori and summations of divergent series*, Communications in Mathematical Physics **227** (2002), no. 3, 421–460.
- 5[GG2] G. Gallavotti, G. Gentile, *Degenerate elliptic resonances*, Communications in Mathematical Physics **257** (2005), 319–362.
- 6[JLZ] À. Jorba, R. de la Llave, M. Zou, *Lindstedt series for lower-dimensional tori*, “Hamiltonian systems with three or more degrees of freedom” (S’Agaró, 1995), 151–167, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 533, Kluwer Acad. Publ., Dordrecht, 1999.

- ⁷[Ne] F. Nevanlinna, *Zur Theorie der Asymptotischen Potenzreihen*, Annales Academiae Scientiarum Fennicae. Series A I. Mathematica **12** (1916), 1–18.
- ⁸[So] A.D. Sokal, *An improvement of Watson's theorem on Borel summability*, Journal of Mathematical Physics **21** (1980), no. 2, 261–263.