

Frictionless thermostats and intensive constants of motion

G. Gallavotti* and E. Presutti[®]

* Fisica-INFN Roma1, [®] Matematica Roma2

September 3, 2009

Abstract: *Thermostats models in space dimension $d = 1, 2, 3$ for nonequilibrium statistical mechanics are considered and it is shown that, in the thermodynamic limit, the evolutions admit infinitely many constants of motion: namely the intensive observables.*

I. THERMOSTATS

Extremal translation invariant DLR distributions, [1] have many constants of motion, among them all intensive observables. In fact the characterizing property of such distributions μ is that they are time invariant, [2], and since the intensive variables are measurable at infinity, [1], they are μ -almost surely constant and therefore invariant at almost all times. The same property holds for all DLR distributions because they are integrals over the extreme ones.

The space-time invariance is essential: however we shall be interested in models in which such invariance properties are not satisfied, and we want also to check that the intensive variables are constant for all times (not just almost all).

The class of models that we shall investigate is when particles of a *test system*, in a container Ω_0 , and ν other particles systems, in containers $\Omega_1, \dots, \Omega_\nu$, interact and define a model of a system in interaction with ν thermostats, if the particles in $\Omega_1, \dots, \Omega_\nu$ can be considered at fixed temperatures T_1, \dots, T_ν .

A representation of the system is in Fig.1:




Fig.1: If $d = 1, 2$ the $1 + \nu$ finite boxes $\Omega_j \cap \Lambda$, $j = 0, \dots, \nu$, are marked C_0, C_1, \dots, C_ν in the first figure and contain N_0, N_1, \dots, N_ν particles, out of the infinitely many particles, with positions and velocities denoted $\mathbf{X}_0, \mathbf{X}_1, \dots, \mathbf{X}_\nu$, and $\dot{\mathbf{X}}_0, \dot{\mathbf{X}}_1, \dots, \dot{\mathbf{X}}_\nu$, respectively, contained in Ω_j , $j \geq 0$. The second figure illustrates the special geometry that will be considered for $d = 1, 2, 3$: here two thermostats, symbolized by the shaded regions, Ω_1, Ω_2 occupy half-spaces adjacent to Ω_0 .

From the point of view of Physics the temperatures in the thermostats are fixed. A natural model, often invoked in the applications, [3], is to imagine the containers Ω_j , $j = 1, \dots, \nu$, as infinite and occupied by particles initially in a Gibbs distribution with given temperatures

and densities $T_1, \delta_1, \dots, T_\nu, \delta_\nu$.

To implement the physical requirement that the thermostats have well defined temperatures and densities the initial data will be imagined to be randomly chosen with a suitable Gibbs distribution

Initial data: *The probability distribution μ_0 for the random choice of initial data will be, if $dx \stackrel{def}{=} \prod_{j=0}^{\nu} \frac{d\mathbf{X}_j d\dot{\mathbf{X}}_j}{N_j!}$, the limit as $\bar{\Lambda} \rightarrow \infty$ of the distributions on the configurations $x \in \mathcal{H}(\bar{\Lambda})$ with $\mathbf{X}_j \in \bar{\Lambda}$ (see Fig.1),*

$$\mu_{0, \bar{\Lambda}}(dx) = \text{const } e^{-H_0(x)} dx \quad (1.1)$$

with $H_0(x) = \sum_{j=0}^{\nu} \beta_j (K_j(\dot{\mathbf{X}}_j) - \lambda_j N_j + U_j(\mathbf{X}_j))$ and $\beta_j \stackrel{def}{=} \frac{1}{k_B T_j} > 0$, $\lambda_j \in \mathbb{R}$, $j > 0$; the values $\beta_0 = \frac{1}{k_B T_0} > 0$, $\lambda_0 \in \mathbb{R}$ will also be fixed.

The values β_0, λ_0 bear no particular physical meaning because the test system is kept finite. Here $\boldsymbol{\lambda} = (\lambda_0, \lambda_1, \dots, \lambda_\nu)$ and $\mathbf{T} = (T_0, T_1, \dots, T_\nu)$ are fixed *chemical potentials* and *temperatures*, and $\bar{\Lambda}$ is a ball centered at the origin and of radius r_0 . The $K_j(\dot{\mathbf{X}}_j), U_j(\mathbf{X}_j)$ are kinetic and potential energies of the particles in Ω_j (see below for the conditions on the potentials).

The distribution μ_0 is interpreted as a Gibbs distribution μ_0 obtained by taking the ‘‘thermodynamic limit’’ $\bar{\Lambda} \rightarrow \infty$. If $p_j(\beta, \lambda; \bar{\Lambda}) \stackrel{def}{=} \frac{1}{\beta |\Omega_j \cap \bar{\Lambda}|} \log Z_j(\beta, \lambda)$ with

$$Z_j(\beta, \lambda) = \sum_{N=0}^{\infty} \int \frac{dx_N}{N!} e^{-\beta(-\lambda N + K_j(x_N) + U_j(x_N))} \quad (1.2)$$

where the integration is over positions and momenta of the N particles in $\bar{\Lambda} \cap \Omega_j$ then we shall say that the thermostats have pressures $p_j(\beta_j, \lambda_j)$, densities δ_j , temperatures T_j , energy densities e_j , and potential energy densities u_j , for $j > 0$, given by equilibrium thermodynamics, *i.e.*:

$$\begin{aligned} p_j(\beta, \lambda) &\stackrel{def}{=} \lim_{\bar{\Lambda} \rightarrow \infty} p_j(\beta_j, \lambda_j, \bar{\Lambda}) \\ \delta_j &= -\frac{\partial p_j(\beta_j, \lambda_j)}{\partial \lambda_j}, \quad k_B T_j = \beta_j^{-1} \\ e_j &= -\frac{\partial \beta_j p_j(\beta_j, \lambda_j)}{\partial \beta_j} - \lambda_j \delta_j, \quad u_j = e_j - \frac{d}{2} \delta_j \beta_j^{-1} \end{aligned} \quad (1.3)$$

which are the relations linking density δ_j , temperature $T_j = (k_B \beta_j)^{-1}$, energy density e_j and potential energy density u_j in a grand canonical ensemble and in absence of phase transitions in correspondence of the parameters (β_j, λ_j) , for $j > 0$.

Remark: (1) notice that the limit defining p_j does not depend on the shape of Ω_j and coincides with the usual definition of pressure in the thermodynamic limit in the sense of Van Hove, [4].

(2) An alternative definition of density, energy density and temperature could be

$$\lim_{n \rightarrow \infty} \left(\frac{N_{j,\Lambda_n}(x)}{|\Lambda_n \cap \Omega_j|}, \frac{U_{j,\Lambda_n}(x)}{|\Lambda_n \cap \Omega_j|}, \frac{K_{j,\Lambda_n}(x)}{N_{j,\Lambda_n}(x)} \right) \quad (1.4)$$

provided the limit exists with μ_0 -probability 1 and is x -independent.

(3) The limits in Eq.(1.4) will exist and be equal to the values in Eq.(1.3) with μ_0 -probability 1 if the hypotheses below hold. This is part of the result of theorem 1.

Hypotheses: *In the geometries of Fig.1 suppose:*

- (1) μ_0 satisfies the DLR equations and that
- (2) the thermostats pressures $p_j(\beta, \lambda)$ are differentiable in β, λ at $\beta_j, \lambda_j, j = 1, \dots, \nu$.

It is essential that the ‘‘macroscopic’’ property of the thermostats, of having given densities and temperatures, remains when the system evolves in time.

Evolution is defined via equations of motion: since we are dealing with infinitely many particles it will be defined by first considering the motion of the particles initially contained in some ball Λ keeping the particles outside Λ fixed. Such motion $x \rightarrow S_t^{(\Lambda)}x$ is called Λ -regularized: then we shall consider the limit as $\Lambda \rightarrow \infty$.

The regularization boxes Λ will be (for simplicity) balls Λ_n centered at the origin O and with radius $2^n r_\varphi$, with r_φ equal to the range of the interparticle potential, and particles will be reflected at the boundary of Λ_n . The limit motion reached as $n \rightarrow \infty$ will define the thermodynamic limit motion.

The Λ_n -regularized equations of motion will be

$$\begin{aligned} m\ddot{\mathbf{X}}_{0i} &= -\partial_i U_0(\mathbf{X}_0) - \sum_{j>0} \partial_i U_{0,j}(\mathbf{X}_0, \mathbf{X}_j) + \boldsymbol{\Phi}_i(\mathbf{X}_0) \\ m\ddot{\mathbf{X}}_{ji} &= -\partial_i U_j(\mathbf{X}_j) - \partial_i U_{0,j}(\mathbf{X}_0, \mathbf{X}_j) \end{aligned} \quad (1.5)$$

(see Fig.1) where:

(1) the first label, $j = 0$ or $j = 1, \dots, \nu$, refers (respectively) to the test system or to a thermostat, while the second indicates the derivatives with respect to the coordinates of the points located in the corresponding container and in the regularization box Λ_n (hence the labels i in the subscripts (j, i) have $N_j d$ values).

(2) The forces $\boldsymbol{\Phi}(\mathbf{X}_0)$ are, positional, *nonconservative*, smooth stirring forces, possibly vanishing; the other forces are conservative and generated by a pair potential φ , with range r_φ , which couples all pairs in the same containers and all pairs of particles one of which is located in Ω_0 and the other in Ω_j (*i.e.* there is *no direct interaction* between the different thermostats).

(3) Furthermore particles are repelled by the boundaries of the containers by a conservative force of potential energy ψ , diverging with the distance r to the walls as $r^{-\alpha}$, for some $\alpha > 0$, and of range $r_\psi \ll r_\varphi$. The potential energies will be $U_j(\mathbf{X}_j)$, $j \geq 0$, and $U_{0,j}(\mathbf{X}_0, \mathbf{X}_j)$, respectively denoting the internal energies of the various systems and the potential energy of interaction between the test system and the thermostats:

$$\begin{aligned} U_j(\mathbf{X}) &= \sum_{q \in \mathbf{X}_j} \psi(q) + \sum_{(q,q') \in \mathbf{X}_j, q \in \Lambda} \varphi(q - q') \\ U_{0,j}(\mathbf{X}_0, \mathbf{X}_j) &= \sum_{q \in \mathbf{X}_0, q' \in \mathbf{X}_j} \varphi(q - q') \end{aligned} \quad (1.6)$$

The potentials φ, ψ have been chosen j -independent for simplicity.

(4) The equations are formally defined also in the phase space \mathcal{H} of the locally finite configurations $x = (\dots, q_i, \dot{q}_i, \dots)_{i=1}^\infty$

$$x = (\mathbf{X}_0, \dot{\mathbf{X}}_0, \mathbf{X}_1, \dot{\mathbf{X}}_1, \dots, \mathbf{X}_n, \dot{\mathbf{X}}_n) = (\mathbf{X}, \dot{\mathbf{X}}) \quad (1.7)$$

with $\mathbf{X}_j \subset \Omega_j$, hence $\mathbf{X} \subset \Omega = \cup_{j=0}^n \Omega_j$, and $\dot{q}_i \in \mathbb{R}^d$; in every ball $\Sigma(r')$ of radius r' and center at the origin O , fall a finite number of points of \mathbf{X} .

Infinite systems are idealizations not uncommon in statistical mechanics. But we take it for granted that they must be considered as limiting cases of large yet finite systems. This leads to several difficulties: one is immediately manifest if one remarks that the equations of motion Eq.(1.5) do not even admit an obvious solution in \mathcal{H} .

Dynamics is well defined with μ_0 -probability 1 because if $d = 1, 2, 3$ the Λ_n -regularized equations with data x admit, with μ_0 -probability 1, a limit $S_t x \stackrel{def}{=} \lim_{\Lambda_n \rightarrow \infty} S_t^{(\Lambda_n)} x$ for all $t > 0$: a precise statement is in theorem 4 below (proved in [5, theorems 6, 7], for $d = 1, 2$, and in [6, Theorem 1] for $d = 1, 2, 3$).

Since the Eq.(1.5) are Newton's equations we shall call the model a *frictionless* thermostats model.

An important question is whether time evolution changes the configuration x into $S_t x$ but keeps the temperatures and densities of the thermostats constant at least with μ_0 -probability 1 and for any finite time. This is part of the more general question whether the spatial average of an intensive observable remains constant in time.

A simple, partial but quantitative, formulation is in terms of the number $N_{j,\Lambda}(S_t x)$ of particles of $S_t x$, of the kinetic energy $K_{j,\Lambda}(S_t x)$ and of the potential energy $U_{j,\Lambda}(S_t x)$ of the configuration $S_t x$ into which x evolves at time t , inside a ball Λ centered at the origin. Consider, then, $\forall j > 0$, the limits (if existent)

$$\lim_{n \rightarrow \infty} \left(\frac{N_{j,\Lambda_n}(S_t x)}{|\Lambda_n \cap \Omega_j|}, \frac{U_{j,\Lambda_n}(S_t x)}{|\Lambda_n \cap \Omega_j|}, \frac{K_{j,\Lambda_n}(S_t x)}{|\Lambda_n \cap \Omega_j|} \right). \quad (1.8)$$

Under the above ‘‘no phase transition’’ assumption on μ_0 we shall prove:

Theorem 1: *The limits in Eq.(1.8) exist with μ_0 -probability 1 for all times and are time independent. The limits will be respectively δ_j, u_j and $\frac{d}{2} \delta_j k_B T_j$ with μ_0 -probability 1, as in Eq.(1.3).*

Remark: This shows that the thermostats keep, in the thermodynamic limit, the same temperature and density that they had in the initial state: a property that has to be required for the model to adhere to the physical intuition behind the empirical notion of thermostats. Hence density and temperature of the thermostats are *constants of motion*. We shall show that more generally many other intensive observables are also constants of motion.

II. INTENSIVE OBSERVABLES

The definition of an h_Γ -particles *intensive observable* is in terms of a smooth function $\Gamma(q_1, \dot{q}_1, \dots, q_h, \dot{q}_h)$ on R^{2dh} vanishing for $h \neq h_\Gamma$ and which is “translation invariant”, and with “short range” r_Γ .

This means that $\Gamma = 0$ if the diameter of $X = (q_1, \dots, q_h)$ exceeds some $r_\Gamma > 0$ and, denoting by $\tau_\xi(X, \dot{X})$ the configuration $(q_1 + \xi, \dot{q}_1, \dots, q_h + \xi, \dot{q}_h)$, it is $\Gamma(\tau_\xi(X, \dot{X})) = \Gamma(X, \dot{X})$, $\forall \xi \in \mathbb{R}^d$.

Given a region W the function G_W of $x = (X, \dot{X})$

$$G_W(x) \stackrel{def}{=} \sum_{Y \subset X \cap W} \Gamma(Y, \dot{Y}) \quad (2.1)$$

defines a “local observable” in $W \subset R^d$ with potential Γ .

We shall say that G_W is an observable of *potential type* if $\Gamma(Y, \dot{Y})$ depends only on Y , while if it depends only on \dot{Y} it will be called of *kinetic type*.

Then, if $V_n \stackrel{def}{=} \Omega_j \cap \Lambda_n$, $|V_n| \stackrel{def}{=} \text{volume}(V_n)$,

Definition 1: The “local average” of Γ on the configuration $x = (X, \dot{X})$ is $|V_n|^{-1} G_{V_n}(x)$. The corresponding “intensive observable” in the j -th thermostat is

$$g(x) \stackrel{def}{=} \lim_{n \rightarrow \infty} \frac{1}{|V_n|} G_{V_n}(x), \quad (2.2)$$

if the limit exists. Furthermore, given μ_0 , define the “intensive fluctuation” of G (in the j -th thermostat)

$$\begin{aligned} \Delta_G(x) &\stackrel{def}{=} \lim_{n \rightarrow \infty} \left(\frac{1}{|V_n|} G_{V_n}(x) - \mu_0 \left(\frac{1}{|V_n|} G_{V_n} \right) \right) \\ &\stackrel{def}{=} \lim_{n \rightarrow \infty} \Delta_{G, V_n}(x), \end{aligned} \quad (2.3)$$

if the limit exists.

Remark: The notation requires keeping in mind that G_{V_n} depends also on j (because $V_n = \Omega_j \cap \Lambda_n$): however for simplicity of notation the labels j on V_n and G_{V_n} will not be marked.

Properties of intensive observables can be derived from various assumptions on the *initial* distributions of the particles in the various regions Ω_j which, we recall, are

distributed independently over $j = 1, \dots, \nu$ and depend on the ν pairs of parameters β_j, λ_j .

The simplest assumption is perhaps the uniqueness of the tangent plane to the graph of the pressure in various directions, which could for instance be insured by the uniqueness of the translation invariant states of our particles system with parameters β_j, λ_j .

Let G be an observable of potential or kinetic type; and suppose that $H_{0,\Lambda,\Gamma}(x) \stackrel{def}{=} H_{0,\Lambda}(x) + \theta G_\Lambda(x)$ is superstable for $|\theta|$ small enough (*i.e.* there exist constants $a > 0, b \geq 0$ such that for all balls Λ it is $H_{0,\Lambda,\Gamma}(x) \geq aN^2/|\Lambda| - bN$ for all configurations $x = (X, \dot{X})$ with N particles and with $X \subset \Lambda$ and $\forall |\theta| \leq \theta_0$ for some $\theta_0 > 0$). We call G an “allowed observable”. For such observables it is possible to define, for $|\theta|$ small, the “pressure”

$$P(\theta) = \lim_{\Lambda \rightarrow \infty} \frac{1}{|\Lambda|} \log \frac{Z_j(\theta)}{Z_j(0)} \quad (2.4)$$

with $Z_j(\theta)$ given by Eq.(1.2) with the energy $\theta G_V(x)$ added in the exponential. It is $P(0) \equiv 0$.

It is important to stress that $P(\theta)$ is, in the geometries in Fig.1 considered here, *independent* of the special geometry considered for the Ω_j as long as the conical containers have d -dimensional shape (*i.e.* they contain balls of arbitrarily large radius).

In this context we can derive the following result:

Theorem 2: Let G be an allowed observable of potential or kinetic type. If $P(\theta)$ is differentiable at $\theta = 0$, then with μ_0 -probability 1 the limit as $|V_n| \rightarrow \infty$ of $\frac{1}{|V_n|} G_{V_n}(S_t x)$ exists μ_0 -almost everywhere and is t -independent.

Remarks: (1) The differentiability assumption of $P(\theta)$ has the meaning of uniqueness of the tangent plane to the graph of the pressure p “in the direction of G ”: such uniqueness is a “generic” property, see [7] for the lattice gas case.

(2) The superstability of $H_{0,\Lambda}(x) + \theta G_\Lambda(x)$ is a very strong condition: it is certainly satisfied if

- (i) $\Gamma(X, \dot{X}) = 1$ for $|X| = 1$ and 0 otherwise, or if
- (ii) $\Gamma(X, \dot{X}) = \frac{1}{2} \dot{q}^2$ for $|X| = 1$ and 0 otherwise, or if
- (iii) $\Gamma(X, \dot{X}) = 0$ unless $X = (q, q')$ and in such case $\Gamma(q, q') = \varphi(q - q')$,

therefore theorem 1 is a corollary of theorem 2.

We also expect that the intensive observables will have very small probability of being appreciably different from their average values, and precisely a probability bounded above by an exponential of the volume $|\Lambda_n|$. This will mean that the observable G satisfies a kind of *large deviations property*:

Theorem 3: Under the assumptions of theorem 2 the μ_0 -probability that the fluctuation $\Delta_{G, \Lambda_n}(S_t x)$ differs

from 0 by more than $\varepsilon > 0$ tends to 0 exponentially fast in $|V_n|$ as $n \rightarrow \infty$, $\forall \varepsilon > 0$.

Remark: The assumptions in theorems 2,3 are satisfied by many observables in the Mayer expansion convergence region in the plane λ_j, β_j , [8]. They are also believed to be satisfied quite generally for observables generated by a potential Γ . In particular they hold generically if Γ is a linear combination of the potentials (i),(ii),(iii) in remark (2) above.

The proof of theorems 2,3 are presented in Sec.IV.

III. TIME EVOLUTION

A quantitative existence theorem of the dynamics can be conveniently formulated in terms of the quantities $v_1 \stackrel{def}{=} \sqrt{2\varphi(0)/m}$, r_φ and $W, \mathcal{N}, v_1, \|x_1\|$ defined as

$$W(x; \xi, R) \stackrel{def}{=} \frac{1}{\varphi(0)} \sum_{q_i \in \mathcal{B}(\xi, R)} \left(\frac{m\dot{q}_i^2}{2} + \frac{1}{2} \sum_{j: j \neq i} \varphi(q_i - q_j) + \psi(q_i) + \varphi(0) \right), \quad (3.1)$$

$\mathcal{N}_\xi(x) \stackrel{def}{=} \text{number particles within } r_\varphi \text{ of } \xi \in \mathbb{R}^d$,

$\|x_i - x'_i\| \stackrel{def}{=} |\dot{q}_i - \dot{q}'_i|/v_1 + |q_i - q'_i|/r_\varphi$

Let $\log_+ z \stackrel{def}{=} \max\{1, \log_2 |z|\}$, $g_\zeta(z) = (\log_+ z)^\zeta$ and

$$\mathcal{E}_\zeta(x) \stackrel{def}{=} \sup_{\xi} \sup_{R > g_\zeta(\xi/r_\varphi)} \frac{W(x; \xi, R)}{R^d}. \quad (3.2)$$

Call \mathcal{H}_ζ the configurations in \mathcal{H} with

$$(1) \quad \mathcal{E}_\zeta(x) < \infty \quad (3.3)$$

$$(2) \quad \frac{N(j, \Lambda_n)}{|\Lambda_n \cap \Omega_j|}, \frac{U(j, \Lambda_n)}{|\Lambda_n \cap \Omega_j|}, \frac{K(j, \Lambda_n)}{|\Lambda_n \cap \Omega_j|} \xrightarrow{n \rightarrow \infty} \delta_j, u_j, \frac{d \delta_j}{2\beta_j}$$

with Λ_n the ball centered at the origin and of radius $2^n r_\varphi$, δ_j, u_j, T_j , given by Eq.(e1.8) if $N(j, \Lambda_n), U(j, \Lambda_n), K(j, \Lambda_n)$ denote the number of particles and their internal potential or kinetic energy in $\Omega_j \cap \Lambda_n$. Each set \mathcal{H}_ζ has μ_0 -probability 1 for $\zeta \geq 1/d$, [2, 9–11]. Then:

Theorem 4: *Let $d \leq 3$, then $\mathcal{H}_{1/d}$ has μ_0 -probability 1 and $S_t x$ exists for μ_0 -almost all $x \in \mathcal{H}_{1/d}$ and $\forall t \geq 0$.*

Given (arbitrarily) a time $\Theta > 0$, if $\mathcal{E} \stackrel{def}{=} \mathcal{E}_{1/d}(x)$, and $|q_i(0)| \leq 2^k r_\varphi$ there are $c = c(\mathcal{E}, \Theta) < \infty, c' = c'(\mathcal{E}, \Theta) > 0$ such that $\forall n \geq k$ and $\forall t \leq \Theta$

$$\begin{aligned} |\dot{q}_i(t)| &\leq c v_1 k^{\frac{1}{2}}, \\ \text{distance}(q_i(t), \partial(\cup_j \Omega_j)) &\geq c' r_\varphi k^{-\frac{1}{\alpha}} \\ \mathcal{N}_\xi(S_t x) &\leq c k^{1/2} \\ \|(S_t x)_i - (S_t^n x)_i\| &\leq e^{-c' 2^{n/2}}, \quad n > k. \end{aligned} \quad (3.4)$$

This is proved in [5, theorem 7] for $d = 2$ and in [6] for $d = 3$ (the latter reference covers also the case $d = 2$ via a somewhat different approach).

Remark that the theorem *does not state* that the second of Eq.(3.3) holds: in [5, 6] it is however proved, in addition to theorem 4, the weaker statement that the \liminf of $\frac{K_{j, \Lambda_n}(S_t x)}{|\Omega_j \cap \Lambda_n|}$ is not smaller than $\frac{1}{2}$ of the corresponding *r.h.s.*; and the same is true for the other two quantities in Eq.(3.3).

A corollary of the main results of this paper will be that the limit relations in Eq.(3.3) will hold for all $t > 0$.

IV. CONSTANTS OF MOTION

Let Γ be an h -points local observable of potential type, $V_n = \Omega_j \cap \Lambda_n$. Under the assumptions of theorem 2 we first show that $\lim_{n \rightarrow \infty} |V_n|^{-1} \langle G_{\Lambda_n} \rangle_{\mu_0} = g$ exists.

Define $P_n(\theta) \stackrel{def}{=} \frac{1}{|V_n|} \log \langle e^{-\theta G_{V_n}} \rangle_{\mu_0}$: this is smooth and convex in θ and its unique derivative at $\theta = 0$ is $g_n \stackrel{def}{=} \frac{1}{|V_n|} \mu_0(G_{V_n})$; therefore, remarking that $P(0) = 0$, it satisfies $P_n(\theta) \geq \theta g_n$.

The limit $P(\theta)$ as $n \rightarrow \infty$ of $P_n(\theta)$ is the same that would be obtained if V_n was replaced by the full ball Λ_n and filled with particles at temperature β_j^{-1} and chemical potential λ_j .

Any convergent subsequence g_{n_i} defines therefore a coefficient g with the property $P(\theta) \geq \theta g$. Hence, by the assumed uniqueness of the tangent to $P(\theta)$ at $\theta = 0$, it follows that g is uniquely determined thus implying that the limit $g \stackrel{def}{=} \lim_{n \rightarrow \infty} g_n$ exists.

Let $g_n = \langle |V_n|^{-1} G_{\Lambda_n} \rangle_{\mu_0}$ and, given $\gamma > 0$, let $\mathcal{X}_{E, \gamma, n}$ to be the set of points in $\mathcal{H}_{1/d}$ with $\mathcal{E}(x) \leq E$, $G_{\Lambda_n}(x) < (g_n + \frac{1}{2}\gamma)|V_n|$ and which, under the evolution, reach in a time $\tau_{\gamma, n}(x) \leq \Theta$ and for the first time, a point of the surface

$$\Sigma_{n, \gamma} \stackrel{def}{=} \{x \mid |V_n|^{-1} G_{\Lambda_n}(x) = (g_n + \gamma)\}. \quad (4.1)$$

If for all E and for all small $\gamma > 0$ it is $\sum_n \mu_0(\mathcal{X}_{E, \gamma, n}) < +\infty$ then it will be $\limsup_{n \rightarrow \infty} |V_n|^{-1} G_{\Lambda_n}(S_t x) \leq g$, with μ_0 -probability 1 (by Borel–Cantelli's estimate); changing Γ into $-\Gamma$ it will follow, again with μ_0 -probability 1, that the \liminf is $\geq g$: notice that the change in sign of Γ is possible by the condition on G to be an “allowed observable”, as introduced before Eq.(2.4).

This remains true if for all small γ there is $\gamma_n \in [\gamma, 2\gamma]$ such that $\sum_n \mu_0(\mathcal{X}_{E, \gamma_n, n}) < +\infty$.

If $x \in \mathcal{X}_{E, \gamma, n}$ the phase space contraction, when phase space volume is measured by μ_0 , within time t is, [5, 6],

$$s(x, t) = \int_0^t \left(\sum_{j \geq 0} \beta_j Q_j(\tau) + \beta_0 L_0(\tau) \right) d\tau \quad (4.2)$$

where $Q_j(t) \stackrel{def}{=} \dot{\mathbf{X}}_j(t) \cdot \mathbf{F}_j$, $L_0(t) \stackrel{def}{=} \dot{\mathbf{X}}_0 \cdot \Phi(\mathbf{X}_0(t))$.

By theorem 4, $L_0(t)$ is uniformly bounded as $n \rightarrow \infty$, for $0 \leq t \leq \Theta$, by the first of Eq.(3.2), by a quantity C (only depending on E, n_0, Θ).

Therefore by a quasi-invariance lemma, [2, 12], [5, Appendix H], $\mu_0(\mathcal{X}_{E, \gamma+\varepsilon, n})$ can be bounded $\forall \varepsilon \in [\gamma, 2\gamma]$ by

$$C \int \mu_0(dx) \frac{|\widehat{G}|}{|V_n|} \delta\left(\frac{G_{\Lambda_n}(x)}{|V_n|} - (g_n + \gamma + \varepsilon)\right) \quad (4.3)$$

where \widehat{G} denotes the time derivative (at $t = 0$) of $G_{\Lambda_n}(S_t x)$ (to be computed via the equations of motion) evaluated on the surface $\Sigma_{n, \gamma+\varepsilon}$, see Eq.(4.1).

Integrating Eq.(4.3) over $d\varepsilon/\gamma$, $\mu_0(\mathcal{X}_{E, n, \gamma_n})$ can be bounded by

$$\frac{C}{\gamma} \int \mu_0(dx) \frac{|\widehat{G}|}{|V_n|} \chi(\gamma \leq \frac{G_{\Lambda_n}(x)}{|V_n|} - g_n \leq 2\gamma), \quad (4.4)$$

with $\widehat{G} = \sum_{X \subset V_n} \sum_{q \in X} \partial_q \Gamma(X) \dot{q}$. By Schwartz' inequality

$$C_2 \gamma^{-1} \mu_0(\{x : \gamma \leq \frac{G_{\Lambda_n}(x)}{|V_n|} - g_n \leq 2\gamma\})^{1/2} \quad (4.5)$$

because from Eq.(2.1) for Γ

$$\mu_0(\widehat{G}^2)^{1/2} \leq C_1 |V_n| \quad (4.6)$$

obtained via superstability bounds, using the Maxwellian distribution for \dot{q} .

The probability in Eq.(4.5) is bounded above by Chebischev inequalities (quadratic or exponential) by both averages

$$I \stackrel{def}{=} \left\langle \frac{(G_{\Lambda_n}(x)/|V_n| - g_n)^2}{\gamma^2} \right\rangle_{\mu_0}, \quad (4.7)$$

$$I_\theta \stackrel{def}{=} \langle e^{\theta(G_{\Lambda_n} - |V_n|(g_n + \gamma))} \rangle_{\mu_0}$$

$\forall \theta \geq 0$. This implies the existence of $\gamma_n \in [\gamma, 2\gamma]$ with:

$$\mu_0(\mathcal{X}_{E, n, \gamma_n}) \leq C_3 \gamma^{-1} J(n), \quad J(n)^2 = I, I_\theta \quad (4.8)$$

Therefore we look for assumptions on the thermostats structure (*i.e.* on $\lambda_j, \beta_j, \varphi$) under which $J(n)$ tends to zero fast enough making $\sum_n \mu_0(\mathcal{X}_{E, n, \gamma_n}) < \infty$. In this case theorem 2 will follow from Borel-Cantelli's lemma and the arbitrariness of γ .

As a consequence of the above bounds, basically following from the uniqueness of the tangent plane in the direction Γ , the proof of theorem 2 can be completed as follows. Fix $\gamma > 0$ and remark that

$$I_\theta = \langle e^{\theta U_{r, v_n}} \rangle_{\mu_0} e^{-\theta(g_n + \gamma)|V_n|} \leq e^{-\theta \gamma |V_n| + \eta(\theta, V_n)} \quad (4.9)$$

Continuing the argument leading to the existence of the limit of g_n , at the beginning of the section, the correction term $\eta(\theta, V_n)$ is bounded, cas follows:

(a) $\frac{1}{|V_n|} \log \langle e^{\theta U_{r, v_n}} \rangle_{\mu_0}$ is $P_n(\theta) - P_n(0)$ (notice: $P_n(0) \equiv 0$) and converges to $P(\theta) - P(0)$ as $V_n \rightarrow \infty$ for $|\theta| \leq \theta_0$, if θ_0 is small enough so that the potential $\varphi + \beta_j^{-1} \theta$ is superstable $\forall |\theta| \leq \theta_0, j = 1, \dots, \nu$. By superstability the limit exists for $|\theta| \leq \theta_0$ and it is a limit of functions $P_n(\theta)$ which are convex for $|\theta| \leq \theta_0$. Hence the limit is uniform: $|P(\theta) - P_n(\theta)| \leq o(|V_n|)$ for $|\theta| \leq \theta_0$,

(b) the g_n in the exponent in Eq.(4.7) has just been shown to be $g_n |V_n| = g |V_n| + o(|V_n|)$, so that $-\theta g_n$ converges to $-\theta g$ with an error $\theta o(|V_n|)$,

(c) $(P(\theta) - P(0) - \theta g) |V_n|$ is (by the uniqueness of the tangent plane) $o(\theta) |V_n|$. Hence

$$\begin{aligned} \eta(\theta, V_n) - \gamma \theta_n |V_n| &\leq -\frac{1}{2} \gamma \theta_n |V_n| \\ &+ \left(-\frac{1}{2} \gamma \theta_n + \frac{o(|V_n|)}{|V_n|} + o(\theta_n) \right) |V_n| \leq -\frac{1}{2} \gamma \theta_n |V_n| \end{aligned} \quad (4.10)$$

and choosing θ_n tending to 0 so slowly that the exponent of the *r.h.s.* of (4.9) tends rapidly to ∞ , for instance if $\theta_n = \max(\frac{1}{\log n}, \frac{1}{4\gamma} \frac{o(|V_n|)}{|V_n|})$, we see that $I_{\theta_n} \xrightarrow{n \rightarrow \infty} 0$ so fast that $\mu_0(\mathcal{X}_{E, n, \gamma_n})$ is summable in n implying theorem 2 and of its special case theorem 1.

Theorem 3 also follows from the existence of the limit for g_n because I_θ yields a summable bound on J , hence on $\mu_0(\Delta_{G, \Lambda_n}^2)$.

Remarks: (1) Uniqueness of the tangent plane can be replaced by assumptions on the decays of correlations in the distribution μ_0 somewhat stronger than just requiring its extremality among the DLR distributions in the geometry in Fig.1.

(2) Sufficient estimates can be formulated as follows: $\rho_j(x_1, \dots, x_n)$ be the n -points correlation function in the j -th container: by superstability $\rho_j \leq C^n$, [9]. If $x = (q, \dot{q})$ and $\xi \in \Omega_j$, extremality of μ_0 , implies, [1, 9], for x_1, \dots, x_n and y_1, \dots, y_m) with positions in Ω_j :

$$\begin{aligned} &|\rho_j(x_1, \dots, x_n, \tau_\xi y_1, \dots, \tau_\xi y_m) \\ &- \rho_j(x_1, \dots, x_n) \rho_j(\tau_\xi y_1, \dots, \tau_\xi y_m)| \xrightarrow{\xi \rightarrow \infty} 0 \end{aligned} \quad (4.11)$$

Assume that Eq.(4.11) holds in the stronger sense that the *l.h.s.* is bounded by $\eta_{R, m, n}(\xi)$ if the positions of x_1, \dots, x_n and y_1, \dots, y_m) can be enclosed in a ball of radius R .

Theorem 5: *If there is a constant $C_{R, m, n} < \infty$ such that $\eta_{R, m, n}(\xi) \leq C_{R, m, n} |\xi|^{-a(R, m, n)}$ with $a(R, m, n) > 0$ and if $\lim_{\Lambda \rightarrow \infty} \frac{1}{|V_n|} \mu_0(G_{V_n}) = g$ exists, then $\lim_{\Lambda \rightarrow \infty}$*

$\frac{1}{|V_n|} \Delta_{G, V_n}(x) = 0$ and $\lim_{\Lambda \rightarrow \infty} \frac{1}{|V_n|} G_{V_n}(S_t x) = g$ with μ_0 probability 1.

Remarks: (1) Thus if μ_0 has a power law cluster property all intensive observables admitting an average value, over space translations, at time 0 are constants of motion.

(2) With the above assumptions we avoid use of the exponential Chebishev inequality and we may thus drop the superstability condition in the definition of the potential Γ . We could actually consider more general observables of the form (in Ω_j)

$$\lim_{n \rightarrow \infty} \frac{1}{|\Omega_j \cap \Lambda_n|} \int_{r \in \mathbb{R}^d: \tau_r \Delta \subset \Omega_j \cap \Lambda_n} \tau_r f(x) dr \quad (4.12)$$

where f is a cylindrical function in Δ (i.e. it does not depend on the particles outside Δ) and τ_r denotes translation by r . If the power law cluster property is satisfied and μ_0 a.s. the limit in (4.12) exists at time 0, then the intensive observables (4.12) are constant of motion under the assumption that f is smooth and grows at most polynomially with the number of particles.

(3) The assumption certainly holds in the cluster expansion convergence region, [4] and [13, Sec.5.9], i.e. high temperature and low density, without extra assumptions.

Proof: Consider the first of Eq.(4.7) and choose $\gamma = \gamma_n = \frac{1}{n}$. The numerator tends to 0 as $|V_n|^{-a(R, n, n)/d}$ if the potential Γ for the observable G_{V_n} vanishes when the diameter of the set $\{x_1, \dots, x_n\}$ exceeds R .

The estimate Eq.(4.8) implies that $\frac{1}{|V_n|} \Delta_{G, V_n}(S_t x)$ tends to 0 with μ_0 -probability 1 for all $t \leq kt_0$ with k integer and $t_0 > 0$ (arbitrarily fixed). Hence if the average of $\frac{1}{|V_n|} \mu_0(G_{V_n})$ exists it exists for all times and has a time-independent value.

V. ENTROPY AND THERMOSTATS

Entropy production per unit time is defined in terms of $Q_j = -\dot{\mathbf{X}}_j \cdot \partial_{\mathbf{X}_j} U_{0,j}(\mathbf{X}_0, \mathbf{X}_j)$, which is the work per unit time, performed by the test system on the j -th thermostat. Since Q_j is interpreted as the *heat* ceded by the system to the thermostats the entropy production in the configuration x is given by $\sigma_0(x) = \sum_{j>0} \beta_j Q_j(x)$.

If the volumes in phase space are measured by the distribution μ_0 this quantity differs from the contraction rate of the phase space volume by $\beta_0(\dot{Q}_0 + L_0) \equiv \beta_0(\dot{K}_0 + \dot{U}_0)$ and $K_0 + U_0$ is *expected* to stay finite uniformly in time. If so the statistics of the long time averages of the phase space contraction rate and of the entropy production rate will coincide (however this is not proved as the theorems above only concern what happens in a *arbitrarily prefixed but finite* time interval).

In other words in the frictionless thermostats model and in the isoenergetic thermostat models, [6], the entropy production can be identified, possibly up to a time derivative of a quantity expected to be uniformly finite in time, with the phase space contraction. Furthermore the entropy production is the same in both models of thermostats if the thermodynamic parameters of the thermostats ($\delta_j, T_j, j > 0$) are the same: this follows from the equivalence theorem between frictionless and isoenergetic thermostats, [6, theorem 1], which states that under such conditions the microscopic motions of the two models starting from the same initial condition remain identical forever with μ_0 -probability 1.

Other thermostats can be considered: for instance the isokinetic thermostats. At a heuristic level analogous conclusions can be reached, [14].

-
- [1] O. Lanford and D. Ruelle. Observables at infinity and states with short range correlations in statistical mechanics. *Communications in Mathematical Physics*, 13:194–215, 1969.
 - [2] C. Marchioro, A. Pellegrinotti, and E. Presutti. Existence of time evolution for ν dimensional statistical mechanics. *Communications in Mathematical Physics*, 40:175–185, 1975.
 - [3] R.P. Feynman and F.L. Vernon. The theory of a general quantum system interacting with a linear dissipative system. *Annals of Physics*, 24:118–173, 1963.
 - [4] D. Ruelle. *Statistical Mechanics*. Benjamin, New York, 1969.
 - [5] G. Gallavotti and E. Presutti. Thermodynamic limit of isoenergetic and hamiltonian thermostats. *arXiv*, 0903.3316:1–9, 2009.
 - [6] G. Gallavotti and E. Presutti. Nonequilibrium, thermostats and thermodynamic limit. *arXiv*, 0905.3150:1–22, 2009.
 - [7] G. Gallavotti and S. Miracle-Solé. Statistical mechanics of lattice systems. *Communications in Mathematical Physics*, 5:317–323, 1967.
 - [8] G. Gallavotti, F. Bonetto, and G. Gentile. *Aspects of the ergodic, qualitative and statistical theory of motion*. Springer Verlag, Berlin, 2004.
 - [9] D. Ruelle. Superstable interactions in classical statistical mechanics. *Communications in Mathematical Physics*, 18:127–159, 1970.
 - [10] J. Fritz and R.L. Dobrushin. Non-equilibrium dynamics of two-dimensional infinite particle systems with a singular interaction. *Communications in Mathematical Physics*, 57:67–81, 1977.
 - [11] E. Caglioti, C. Marchioro, and M. Pulvirenti. Non-equilibrium dynamics of three-dimensional infinite particle systems. *Communications in Mathematical Physics*, 215:25–43, 2000.
 - [12] Ya. G. Sinai. Construction of dynamics in one-dimensional systems of statistical mechanics. *Theoretical and Mathematical Physics*, 11:487–494, 1972.
 - [13] G. Gallavotti. *Statistical Mechanics. A short treatise*. Springer Verlag, Berlin, 2000.
 - [14] G. Gallavotti. On thermostats: Isokinetic or Hamiltonian? finite or infinite? *Chaos*, 19:013101 (+7), 2008.